

DIFFERENTIAL DIAGNOSIS OF GYNECOLOGIC ORGAN-RELATED DISEASES IN WOMEN PRESENTING WITH ASCITES

Ming-Huei Cheng^{1,2}, Ming-Shyen Yen¹, Kuan-Chong Chao¹, Bor-Ching Sheu³, Peng-Hui Wang^{1,2*}

¹Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, ²Institute of Clinical Medicine, National Yang-Ming University School of Medicine, and ³Department of Obstetrics and Gynecology, National Taiwan University Hospital and National Taiwan University, Taipei, Taiwan.

SUMMARY

Ascites is a pathologic accumulation of fluid within the peritoneal cavity, and usually develops as a result of liver disease, congestive heart failure or nephrotic syndrome. Ascites is also a common manifestation of some gynecologic diseases. It is important that health care workers consider gynecologic problems among the potential differential diagnoses in patients presenting with ascites. Various kinds of ovarian diseases, such as epithelial ovarian cancer, benign ovarian fibroma, stromal hyperplasia, ovarian hyperstimulation syndrome, primary peritoneal serous carcinoma, endometriosis and peritoneal tuberculosis, should be kept in mind when women are found to have ascites. [*Taiwan J Obstet Gynecol* 2008;47(4):384–390]

Key Words: ascites, endometriosis, epithelial ovarian cancer, ovarian hyperstimulation syndrome, peritonitis tuberculosis

Introduction

Ascites is a term that describes a pathologic accumulation of fluid within the peritoneal cavity. It can develop as a result of liver disease, congestive heart failure or nephrotic syndrome, and is not an uncommon complication accompanying some gynecologic diseases. It is, therefore, important that health care workers consider gynecologic problems among the potential differential diagnoses in patients with ascites. The following review explores the gynecologic diseases that can result in ascites.

Ovarian Malignancy

Epithelial ovarian cancer is possibly the most common type of cancer leading to malignant ascites [1–4]. In a 2-year, single institute, retrospective analysis, 37.7% of

patients with ovarian cancer developed ascites, accounting for a quarter of the cases of malignant ascites [1]. Paracentesis is commonly used to evaluate ascites and to make a differential diagnosis. Around 70% of patients with ovarian cancer demonstrate cytologic evidence of ascites [5]. Nevertheless, in some cases where an ovarian malignancy is suspected, paracentesis may be withheld because of the risk of abdominal wall metastases developing at the paracentesis entry sites, especially since the occurrence of this phenomenon, together with the presence of ascites, is highly associated with advanced tumor stage [6,7]. The reported incidence of tumor implantations at trocar or puncture sites after invasive diagnostic procedures varies widely, ranging from 1.2–13.7% [6–9]. The mechanism behind puncture site metastasis is not well understood, but it is likely to be related to direct implantation following surgical trauma [10,11]. The concept of standard therapy for advanced ovarian cancer using primary debulking surgery followed by chemotherapy has recently been challenged by an alternative option using neo-adjuvant chemotherapy followed by interval debulking surgery [12–14]. This would increase the possibility for performing diagnostic paracentesis before getting a definite diagnosis. Although the presence of



*Correspondence to: Dr Peng-Hui Wang, Department of Obstetrics and Gynecology, Taipei Veterans General Hospital and National Yang-Ming University School of Medicine, 201, Section 2, Shih-Pai Road, Taipei 112, Taiwan.
E-mail: phwang@vghtpe.gov.tw
Accepted: August 27, 2008

abdominal wall metastasis does not seem to be significantly correlated with survival [7], the potential danger of performing diagnostic paracentesis should always be kept in mind.

The presence of malignant ascites is usually indicative of a poor prognosis; however, malignant ascites of ovarian origin has a relatively higher survival rate than that of other origins [1–4]. The management of malignant ascites caused by operable ovarian cancer differs from that of other cancers. The mainstay of therapy for ascites accompanying ovarian cancer, unlike other cancers, is attempted removal of all detectable tumors by debulking surgery, followed by adjuvant chemotherapy. The ascites will subside if the ovarian cancer is responsive to chemotherapy. However, for those with malignant ascites caused by recurrent or refractory ovarian cancer, palliative treatment by serial paracentesis or drainage is a possible option to provide symptomatic relief [15–17]. Infectious morbidity is a major concern associated with invasive procedures. In addition, the removal of large volumes of ascitic fluid may result in protein loss and hypovolemia, with subsequent circulatory problems. Although diuretics and salt restriction have been empirically used in the treatment of ascites caused by different kinds of medical illnesses, the effectiveness of these approaches for malignant ascites has yet to be proven. Treatment of the causative malignant cells by intraperitoneal cytotoxic chemotherapy provides another possible approach for controlling malignant ascites. Various antineoplastic agents, including cisplatin, carboplatin, 5-fluorouracil and leucovorin, as well as metalloproteinase inhibitors, have been used to inhibit ascites formation. Intraperitoneal agents that block vascular endothelial growth factor have also been found to be a promising therapy for ascites related to carcinoma in both animal models and human clinical trials [18,19]. Some experimental strategies, such as the use of radiolabeled antibodies and immunotherapy, have been reported to be alternative options for ovarian cancer patients with malignant ascites [20].

Pelvic or Peritoneal Tuberculosis

Tuberculosis is a mycobacterial infection that most commonly manifests as pulmonary disease, but may also affect extrapulmonary sites [21–23], such as the abdominopelvic region. It usually affects women of reproductive age [24]. The clinical presentation of pelvic tuberculosis is non-characteristic, and includes vague pelvic pain, abdominal distension, weight loss or fever. Infertility is a relatively frequent complaint in female patients with pelvic tuberculosis [25]. On diagnostic

imaging, pelvic tuberculosis may appear as ascites, peritoneal nodules, omental or peritoneal thickening, or adnexal masses. The clinical manifestations and imaging features can mimic disseminated carcinomatosis [26–36]. However, there are some subtle ultrasonographic differences that can help to distinguish between pelvic tuberculosis and ovarian cancer: (1) tuberculous ovarian masses are usually almost unenlarged; (2) the septum inside the tuberculous ovarian mass is usually composed of multiple, delicate, incomplete septa, rather than of thick bands as found in ovarian malignancy; and (3) the blood vessels supplying the tuberculous implants are present in high impedance, which reflects vascular resistance, calculated after measuring peak systolic (S) and peak diastolic (D) velocities of blood flow by colored Doppler ultrasound using the formula: resistance index = $(S - D)/S$ [37].

Ascites is a common presentation of pelvic tuberculosis. However, the diagnostic value of paracentesis in ascitic fluid analysis is still inconclusive. The sensitivity of direct preparations of ascitic fluid, for either acid-fast stain or tuberculosis culture, is relatively low [38]. Moreover, the long turnaround time required for culture of the microorganism makes this method impractical. Currently, ascitic fluid analysis by polymerase chain reaction for mycobacteria or for elevated adenine deaminase levels is considered to be a reliable diagnostic technique [39,40]. Blind percutaneous peritoneal biopsy was first advocated 40 years ago as the procedure of choice in cases of early tuberculous peritonitis [41], with varying reported rate of success [42]. Peritoneal biopsy was performed at the left lower quadrant of the abdomen, adjacent to the rectus muscle, using an Abrams needle or Cope needle, under local anesthesia. The limitations include complications such as bowel perforation and even mortality, and the technique is contraindicated in the absence of ascites. However, these limitations could be overcome by using ultrasound- or computed tomography-guided needle biopsy, which is deserving of further study [43]. Even though pelvic tuberculosis is usually secondary to previous tuberculosis infection, the tuberculin skin test and chest roentgenograms do not help to distinguish patients with pelvic tuberculosis [44,45]. Most patients in the literature were diagnosed by exploratory laparotomy and by confirmatory histopathology.

With significant improvements in laparoscopic techniques, laparoscopy has become an effective diagnostic tool to confirm uncertain abdominal conditions [46], and provides an alternative to traditional laparotomy [47] and a better choice than exploratory laparotomy [48]. The characteristic laparoscopic appearance includes free ascites with multiple yellowish-white

nodules of tubercles on the visceral and parietal peritoneum, peritoneal or visceral adhesion, and occasionally inflamed hemorrhagic areas on the peritoneum [42]. The results of a large case series that investigated laparoscopic evaluation (combined with peritoneal biopsy) of ascites of unknown origin found that a definite diagnosis was established in 80–97% of cases [42, 49–51]. Patients diagnosed with pelvic tuberculosis had a satisfactory response to anti-tuberculosis drugs and ascites generally resolved after anti-tuberculosis treatment [29,32].

Ovarian Hyperstimulation Syndrome (OHSS)

Ascites is a clinical manifestation of OHSS. OHSS is an iatrogenic complication of ovarian stimulation for assisted reproductive techniques [52,53]. The pathogenesis of OHSS may be associated with an exaggerated response to gonadotropin, resulting in increased vascular permeability via the activity of vascular endothelial growth factor, tumor necrosis factor, and interleukin. The syndrome is characterized by cystic enlargement of the ovaries and fluid leakage from the blood vessels into the interstitial space [52]. The clinical presentation of OHSS varies considerably, ranging from mild distension, discomfort and nausea, to dyspnea, oliguria, increased hematocrit, thromboembolism, or even abnormal creatinine clearance, liver dysfunction or respiratory distress. The diagnosis is usually established by a history of assisted reproductive technique, clinical appearance, and specific ultrasound findings. Examination of the abdomen should be deferred to prevent rupture of enlarged ovaries and subsequent hemorrhage. The most recent classification of OHSS was introduced in 1999 [54] and differs from previous systems in that a mild degree of OHSS is omitted from the classification, while severe OHSS is classified into three grades, depending on general clinical symptoms and laboratory parameters. The most effective treatment for OHSS is prevention [55]. For OHSS-associated ascites, ultrasound-guided paracentesis has been shown to result in the relief of abdominal distension, improvement of dyspnea, increased diuresis, and decreased hematocrit [56–59]. Concurrent intravenous rehydration and colloid solution supplement are needed to maintain euolemia [60]. Only a few pharmacologic interventions are available to ameliorate extravascular leakage; diuretics play no role [61], but dopamine treatment and dopamine prodrugs have been used to reduce ascites [62,63]. A prospective, randomized, double-blind study in oocyte donors at risk of developing

OHSS found that administration of cabergoline, a potent dopamine agonist, significantly reduced the occurrence of ascites [64].

Meigs Syndrome, Pseudo-Meigs Syndrome, and Pseudo-pseudo Meigs Syndrome

Meigs syndrome is defined as a triad of benign ovarian tumors accompanied by ascites and pleural effusion [65]. The ovarian tumor may be a fibroma, thecoma or granulosa tumor, of which ovarian fibromas are the most common cause of Meigs syndrome. Ovarian fibromas constitute less than 5% of all ovarian tumors, and only 1% of diagnosed cases have been associated with the clinical picture of Meigs syndrome [66,67]. Meigs syndrome is a rare clinical entity. Pseudo-Meigs syndrome is a variant with an identical clinical constellation, but without the original benign tumor cell types described by Meigs [68]. The benign tumors include those of the fallopian tubes and uterus, mature teratomas, struma ovarii, and ovarian leiomyomas. The distinction between Meigs syndrome and pseudo-Meigs syndrome is mostly academic, because the therapeutic strategy is the same in both cases [69]. The pathophysiology of ascites in Meigs syndrome or pseudo-Meigs syndrome remains enigmatic. Some have speculated that the accumulation of peritoneal fluid is caused by irritation of the peritoneal surfaces or by direct pressure on the surrounding lymphatics or vessels by the solid ovarian tumor [70–72]. Other proposed mechanisms suggest that the ascites is caused by fluid secretion from the ovarian tumor, or as a result of increased capillary permeability via release of mediators from the tumor [65]. The ascites usually disappears after resection of the tumor, and the prognosis is favorable.

In contrast to Meigs syndrome or pseudo-Meigs syndrome, pseudo-pseudo Meigs syndrome is a term that describes patients with ascites, pleural effusion, and enlarged ovaries secondary to systemic lupus erythematosus [73]. The ovaries in pseudo-pseudo Meigs syndrome may be enlarged, but without evidence of a tumor. Ascites in pseudo-pseudo Meigs syndrome is usually exudative, and may be associated with activated mesothelial cells in systemic lupus erythematosus [73,74].

Endometriosis

Endometriosis is a relatively common gynecologic disease, defined as the ectopic presence of endometrial epithelium and stroma outside the uterine cavity. Common clinical manifestations of endometriosis

include pelvic pain, pelvic tenderness, dysmenorrhea, dyspareunia, and a history of infertility. The fact that endometriosis can rarely be associated with ascites is not well known, even amongst practicing gynecologists. Fewer than 50 such cases have been reported since the first case was described in 1954 [75–91]. The vast majority of patients were nulliparous non-Caucasian women of reproductive age [76]. The ascites is mostly massive and may be concurrent with pleural effusion. Because endometriosis-associated ascites usually presents with a pelvic mass, poor appetite, weight loss and elevated tumor markers for ovarian neoplasms, it is commonly mistaken for ascites caused by ovarian neoplasms. Although cytologic examination via paracentesis has been used successfully to detect endometriosis in effusion [92], most patients were diagnosed by operative assessment and histologic confirmation. The pathogenesis of ascites associated with endometriosis is speculative. Ascites can form as a result of the rupture of endometriomas [89,93] or, rarely, from hepatic involvement by cystic endometriosis [86]. The hypothesis that endometrial cells shedding into the peritoneal cavity may irritate the peritoneum, thereby leading to ascites, is also widely accepted [94]. The definitive therapy for endometriosis-related ascites is, therefore, to eliminate ovarian function by oophorectomy. Medical treatment using either a progestational agent or gonadotropin-releasing hormone (GnRH) agonist is less reliable than surgical castration, and may lead to a failed response and a recurrence of ascites. However, GnRH agonists have been used successfully to manage various kinds of endometriosis and its related diseases [95–100].

Conclusion

This review indicates the need for clinicians to examine possible gynecologic sources in female patients presenting with ascites (Table). The initial evaluation of a female patient with ascites should rely on a detailed history and physical examination. Paracentesis provides

a rapid and simple means of analyzing ascitic fluid, but its effectiveness in determining the underlying cause is limited in patients with ascites resulting from gynecologic disorders. The treatment of ascites depends on the cause of fluid retention. Most cases of gynecologic disease-associated ascites can be resolved following an accurate diagnosis and appropriate therapy.

Acknowledgments

This article was supported, in part, by grants (96VN-008 and 97VN-012) from Taipei Veterans General Hospital-National Taiwan University Hospital Joint Research Program.

References

1. Ayantunde AA, Parsons SL. Pattern and prognostic factors in patients with malignant ascites: a retrospective study. *Ann Oncol* 2007;18:945–9.
2. Mackey JR, Venner PM. Malignant ascites: demographics, therapeutic efficacy and predictors of survival. *Can J Oncol* 1996;6:474–80.
3. Parsons SL, Lang MW, Steele RJC. Malignant ascites: a 2-year review from a teaching hospital. *Eur J Surg Oncol* 1996;22:237–9.
4. Wilailak S, Linasmita V, Srivannaboon S. Malignant ascites in female patients: a seven-year review. *J Med Assoc Thai* 1999; 82:15–9.
5. Hirasawa T, Yasuda M, Muramatsu T, et al. Cytologic study of ascites and the endometrium in ovarian carcinoma. Clinical significance. *Acta Cytol* 1997;41:1451–5.
6. Kruitwagen RF, Swinkels BM, Keyser KG, Doesburg WH, Schijf CP. Incidence and effect on survival of abdominal wall metastases at trocar or puncture sites following laparoscopy or paracentesis in women with ovarian cancer. *Gynecol Oncol* 1996;60:233–7.
7. Nagarsheth NP, Rahaman J, Cohen CJ, Gretz H, Nezhat F. The incidence of port-site metastases in gynecologic cancers. *JSL* 2004;8:133–9.
8. Aqua KA, Surwit EA, Childers JM. Abdominal-wall tumor implantation following laparoscopic for malignant conditions. *J Am Assoc Gynecol Laparosc* 1994;1:S1–2.
9. Childers JM, Aqua KA, Surwit EA, Hallum AV, Hatch KD. Abdominal-wall tumor implantation after laparoscopy for malignant conditions. *Obstet Gynecol* 1994;84:765–9.
10. Wang PH, Yen MS, Yuan CC, CHao KC, Ng HT, Lee WL, Chao HT. Port site metastasis after laparoscopic-assisted vaginal hysterectomy for endometrial cancer: possible mechanisms and prevention. *Gynecol Oncol* 1997;66:151–5.
11. Wang PH, Yuan CC, Lin G, Ng HT, Chao HT. Risk factors contributing to early occurrence of port site metastases of laparoscopic surgery for malignancy. *Gynecol Oncol* 1999; 72:38–44.

Table. Characteristics of ascites resulting from various kinds of gynecologic diseases

Disease	Characteristics of ascites
Epithelial ovarian cancer	Exudate
Meigs syndrome	Most transudate, but few hemorrhagic exudate
Tuberculosis	Exudate
Ovarian hyperstimulation syndrome	Exudate
Endometriosis	Exudate

12. Schmeler KM, Sun CC, Bodurka DC, et al. Neoadjuvant chemotherapy for low-grade serous carcinoma of the ovary or peritoneum. *Gynecol Oncol* 2008;108:510-4.
13. Morrison J, Swanton A, Collins S, Kehoe S. Chemotherapy versus surgery for initial treatment in advanced ovarian epithelial cancer. *Cochrane Database Syst Rev* 2007;(4):CD005343.
14. Hou JY, Kelly MG, Yu H, McAlpine JN, Azodi M, Rutherford TJ, Schwartz PE. Neoadjuvant chemotherapy lessens surgical morbidity in advanced ovarian cancer and leads to improved survival in stage IV disease. *Gynecol Oncol* 2007;105:211-7.
15. Brooks RA, Herzog TJ. Long-term semi-permanent catheter use for the palliation of malignant ascites. *Gynecol Oncol* 2006;101:360-2.
16. Gotlieb WH, Feldman B, Feldman-Moran O, et al. Intraperitoneal pressures and clinical parameters of total paracentesis for palliation of symptomatic ascites in ovarian cancer. *Gynecol Oncol* 1998;71:381-5.
17. Easson A, Bezjak A, Ross S, Sam F, Newman A, Bana N, Wright JG. Changes in symptoms after paracentesis for symptomatic malignant ascites. *J Clin Oncol* 2005;23:6071.
18. Hamilton CA, Maxwell GL, Chernofsky MR, Bernstein SA, Farley JH, Rose GS. Intraperitoneal bevacizumab for the palliation of malignant ascites in refractory ovarian cancer. *Gynecol Oncol* 2008;doi:10.1016/j.ygyno.2008.04.028.
19. Numnum TM, Rocconi RP, Whitworth J, Barnes MN. The use of bevacizumab to palliate symptomatic ascites in patients with refractory ovarian carcinoma. *Gynecol Oncol* 2006;102:425-8.
20. Burges A, Wimberger P, Kümper C, et al. Effective relief of malignant ascites in patients with advanced ovarian cancer by a trifunctional anti-EpCAM x anti-CD3 antibody: a phase I/II study. *Clin Cancer Res* 2007;13:3899-905.
21. Wang PH, Yuan CC. Tuberculous peritonitis. *Am J Obstet Gynecol* 2001;184:250-1.
22. Wang PH. Tuberculous peritonitis should be always considered as a differential diagnosis in abdominal carcinomatosis. *Taiwan J Obstet Gynecol* 2004;43:182-4.
23. Huang YJ, Wei LH, Hsieh CY. Clinical presentation of pelvic tuberculosis imitating ovarian malignancy. *Taiwan J Obstet Gynecol* 2004;43:29-34.
24. Sutherland AM. The changing pattern of tuberculosis of the female genital tract: a thirty-year survey. *Arch Gynecol* 1983;234:95-101.
25. Sharma JB, Roy KK, Pushparaj M, Kumar S, Malhotra N, Mittal S. Laparoscopic findings in female genital tuberculosis. *Arch Gynecol Obstet* 2008;278:359-64.
26. Gürgen T, Zeyneloğlu H, Urman B, Develioğlu O, Yalali H. Pelvic-peritoneal tuberculosis with elevated serum and peritoneal fluid Ca-125 levels. A report of two cases. *Gynecol Obstet Invest* 1993;35:60-1.
27. Odejinmi F, Annan HG, Hussein SY. Tuberculosis, the great mimic again? A report of two cases of pelvic tuberculosis initially suspected to be advanced ovarian carcinoma. *J Obstet Gynaecol* 2000;20:544-5.
28. Sinha P, Johnson AN, Chidamberan-Pillai S. Pelvic tuberculosis: an uncommon gynaecological problem presenting as ovarian mass. *BJOG* 2000;107:139-40.
29. Bilgin T, Karabay A, Dolar E, Develioğlu OH. Peritoneal tuberculosis with pelvic abdominal mass, ascites and elevated CA 125 mimicking advanced ovarian carcinoma: a series of 10 cases. *Int J Gynecol Cancer* 2001;11:290-4.
30. Koc S, Beydilli G, Tulunay G, et al. Peritoneal tuberculosis mimicking advanced ovarian cancer: a retrospective review of 22 cases. *Gynecol Oncol* 2006;103:565-9.
31. Chow TW, Lim BK, Vallipuram S. The masquerades of female pelvic tuberculosis: case reports and review of literature on clinical presentations and diagnosis. *J Obstet Gynaecol Res* 2002;28:203-10.
32. İlhan AH, Durmusoglu F. Case report of a pelvic-peritoneal tuberculosis presenting as an adnexial mass and mimicking ovarian cancer, and a review of the literature. *Infect Dis Obstet Gynecol* 2004;12:87-9.
33. Mahdavi A, Malviya VK, Herschman BR. Peritoneal tuberculosis disguised as ovarian cancer: an emerging clinical challenge. *Gynecol Oncol* 2002;84:167-70.
34. Wang YC, Lu JJ, Chen CH, Peng YJ, Yu MH. Peritoneal tuberculosis mimicking ovarian cancer can be diagnosed by polymerase chain reaction: a case report. *Gynecol Oncol* 2005;97:961-3.
35. Manidakis LG, Angelakis E, Sifakis S, Stefanaki P, Kalogeraki A, Manidaki A, Koumantakis E. Genital tuberculosis can present as disseminated ovarian carcinoma with ascites and raised Ca-125: a case report. *Gynecol Obstet Invest* 2001;51:277-9.
36. Piura B, Rabinovich A, Leron E, Yanai-Inbar I, Mazor M. Peritoneal tuberculosis mimicking ovarian carcinoma with ascites and elevated serum CA-125: case report and review of literature. *Eur J Gynaecol Oncol* 2002;23:120-2.
37. Wang PH, Yuan CC, Yu KJ, Chao HT. High resistance index of Doppler ultrasound in tuberculous peritonitis presenting as abdominal carcinomatosis. *Chin Med J (Taipei)* 1998;61:175-9.
38. Uygur-Bayramicli O, Dabak G, Dabak R. A clinical dilemma: abdominal tuberculosis. *World J Gastroenterol* 2003;9:1098-101.
39. Ginesu F, Pirina P, Sechi LA, et al. Microbiological diagnosis of tuberculosis: a comparison of old and new methods. *J Chemother* 1998;10:295-300.
40. Marshall JB. Tuberculosis of the gastrointestinal tract and peritoneum. *Am J Gastroenterol* 1993;88:989-99.
41. Donohoe RF, Shnider BI, Gorman J. Needle biopsy of the peritoneum; a preliminary report. *AMA Arch Intern Med* 1959;103:739-45.
42. Chow KM, Chow VC, Szeto CC. Indication for peritoneal biopsy in tuberculous peritonitis. *Am J Surg* 2003;185:567-73.
43. Mizunoe S, Morinaga R, Umeki K, et al. A case of tuberculous peritonitis diagnosed by ultrasonography-guided peritoneal biopsy. *Kansenshogaku Zasshi* 2000;74:589-93. [In Japanese]
44. Ylinen O. Genital tuberculosis in women: clinical experiences with 348 proved cases. *Acta Obstet Gynecol Scand Suppl* 1961;40(Suppl 2):1-213.
45. Bhansali SK. Abdominal tuberculosis. *Am J Gastroenterol* 1977;67:324-7.

46. Wang PH, Yuan CC, Chao HT, Shu LP, Lai CR. Isolated tubal torsion managed laparoscopically. *J Am Assoc Gynecol Laparosc* 2000;7:423-7.
47. Wang PH, Liu WM, Fuh JL, Chao HT, Yuan CC, Chao KC. Laparoscopic uterine vessel occlusion in the treatment of women with symptomatic uterine myomas with and without adding laparoscopic myomectomy—4-year results. *J Minim Invasive Gynecol* 2008;15:712-8.
48. Wang PH, Lee WL, Yuan CC, Chin TW. Postpartum sterilization: the choice of laparoscopy or minilaparotomy? *Eur J Obstet Gynecol Reprod Biol* 2008;139:116-7.
49. Panoskaltis TA, Moore DA, Haidopoulos DA, McIndoe AG. Tuberculous peritonitis: part of the differential diagnosis in ovarian cancer. *Am J Obstet Gynecol* 2000;182:740-2.
50. Inadomi JM, Kapur S, Kinkhabwala M, Cello JP. The laparoscopic evaluation of ascites. *Gastrointest Endosc Clin N Am* 2001;11:79-91.
51. Sandikci MU, Colakoglu S, Ergun Y, Unal S, Akkiz H, Sandikci S, Zorludemir S. Presentation and role of peritoneoscopy in the diagnosis of tuberculous peritonitis. *J Gastroenterol Hepatol* 1992;7:298-301.
52. Beerendonk CC, van Dop PA, Braat DD, Merkus JM. Ovarian hyperstimulation syndrome: facts and fallacies. *Obstet Gynecol Surv* 1998;53:439-49.
53. Budev MM, Arroliga AC, Falcone T. Ovarian hyperstimulation syndrome. *Crit Care Med* 2005;33(Suppl 10):S301-6.
54. Rizk B, Aboulghar MA. Classification, pathophysiology and management of ovarian hyperstimulation syndrome. In: Brinsden P, ed. *In Vitro Fertilization and Assisted Reproduction*. New York: Parthenon Publishing Group, 1999:131-55.
55. Aboulghar MA, Mansour RT. Ovarian hyperstimulation syndrome: classifications and critical analysis of preventive measures. *Hum Reprod Update* 2003;9:275-89.
56. Fluker MR, Copeland JE, Yuzpe AA. An ounce of prevention: outpatient management of the ovarian hyperstimulation syndrome. *Fertil Steril* 2000;73:821-4.
57. Levin I, Almog B, Avni A, Baram A, Lessing JB, Gamzu R. Effect of paracentesis of ascitic fluids on urinary output and blood indices in patients with severe ovarian hyperstimulation syndrome. *Fertil Steril* 2002;77:986-8.
58. Maslovitz S, Jaffa A, Eytan O, Wolman I, Many A, Lessing JB, Gamzu R. Renal blood flow alteration after paracentesis in women with ovarian hyperstimulation. *Obstet Gynecol* 2004;104:321-6.
59. Delvigne A, Rozenberg S. Review of clinical course and treatment of ovarian hyperstimulation syndrome (OHSS). *Hum Reprod Update* 2003;9:77-96.
60. Chen CD, Wu MY, Yang JH, Chen SU, Ho HN, Yang YS. Intravenous albumin does not prevent the development of severe ovarian hyperstimulation syndrome. *Fertil Steril* 1997;68:287-91.
61. Chen SU, Chen CD, Yang YS. Ovarian hyperstimulation syndrome (OHSS): new strategies of prevention and treatment. *J Formos Med Assoc* 2008;107:509-12.
62. Alvarez C, Alonso-Muriel I, García G, Crespo J, Bellver J, Simón C, Pellicer A. Implantation is apparently unaffected by the dopamine agonist cabergoline when administered to prevent ovarian hyperstimulation syndrome in women undergoing assisted reproduction treatment: a pilot study. *Hum Reprod* 2007;22:3210-4.
63. Tsunoda T, Shibahara H, Hirano Y, et al. Treatment for ovarian hyperstimulation syndrome using an oral dopamine prodrug, docarpamine. *Gynecol Endocrinol* 2003;17:281-6.
64. Alvarez C, Marti-Bonmati L, Novella-Maestre E, et al. Dopamine agonist cabergoline reduces hemoconcentration and ascites in hyperstimulated women undergoing assisted reproduction. *J Clin Endocrinol Metab* 2007;92:2931-7.
65. Meigs JV. Fibroma of the ovary with ascites and hydrothorax; Meigs' syndrome. *Am J Obstet Gynecol* 1954;67:962-85.
66. Dockerty MB, Masson JC. Ovarian fibromas: a clinical and pathologic study of two hundred and eighty-three cases. *Am J Obstet Gynecol* 1944;47:741-52.
67. Sivanesaratnam V, Dutta R, Jayalakshmi P. Ovarian fibroma—clinical and histopathological characteristics. *Int J Gynaecol Obstet* 1990;33:243-7.
68. Meigs JV. Pelvic tumors other than fibromas of the ovary with ascites and hydrothorax. *Obstet Gynecol* 1954;3:471-86.
69. Kazanov L, Ander DS, Enriquez E, Jaggi FM. Pseudo-Meigs' syndrome. *Am J Emerg Med* 1998;16:404-5.
70. Stratan D, Socolov R. Meigs' syndrome. A case report and review of literature. *Rev Med Chir Soc Med Nat Lasi* 1991;96:30-2.
71. Terada S, Suzuki N, Uchide K, Akasofu K. Uterine leiomyoma associated with ascites and hydrothorax. *Gynecol Obstet Invest* 1992;33:54-8.
72. Amr SS, Hassan AA. Struma ovarii with pseudo-Meigs' syndrome: report of a case and review of the literature. *Eur J Obstet Gynecol Reprod Biol* 1994;55:205-8.
73. Schmitt R, Weichert W, Schneider W, Luft FC, Kettritz R. Pseudo-pseudo Meigs' syndrome. *Lancet* 2005;366:1672.
74. Ural UM, Kilic A, Gungor T, Ozdal B, Mollamahmutoglu L. Tjalma's or pseudo-pseudo-Meigs' syndrome: a case report. *Clin Exp Dermatol* 2008;33:363-4.
75. Brews A. Endometriosis including endometriosis of the diaphragm and Meigs' syndrome. *Proc R Soc Med* 1951;47:461.
76. Spitzer M, Benjamin F. Ascites due to endometriosis. *Obstet Gynecol Surv* 1995;50:628-31.
77. Myers TJ, Arena B, Granai CO. Pelvic endometriosis mimicking advanced ovarian cancer: presentation with pleural effusion, ascites, and elevated serum CA 125 level. *Am J Obstet Gynecol* 1995;173:966-7.
78. Flanagan KL, Barnes NC. Pleural fluid accumulation due to intra-abdominal endometriosis: a case report and review of the literature. *Thorax* 1996;51:1062-3.
79. Muneyyirci-Delale O, Neil G, Serur E, Gordon D, Maiman M, Sedlis A. Endometriosis with massive ascites. *Gynecol Oncol* 1998;69:42-6.
80. Samora-Mata J, Feste JR. Endometriosis ascites: a case report. *JSLs* 1999;3:229-31.
81. Fletcher H, McFarlane M, Shirley SE, Clarke WF, Lyon K. Massive ascites secondary to severe endometriosis. *West Indian Med J* 1999;48:158-9.
82. El Khalil T, Mourad FH, Barada K, Uthman S. Massive hemorrhagic ascites secondary to endometriosis. *J Clin Gastroenterol* 1999;29:344-5.

83. Crescini C, von Wunster S, Spazzini D. Ascites and pleural and pericardial effusion after operative laparoscopy. *J Am Assoc Gynecol Laparosc* 2000;7:155-7.
84. Bhojawala J, Heller DS, Cracchiolo B, Sama J. Endometriosis presenting as bloody pleural effusion and ascites—report of a case and review of the literature. *Arch Gynecol Obstet* 2000;264:39-41.
85. Dias CC, Andrade JM, Ferriani RA, Villanova MG, Meirelles RS. Hemorrhagic ascites associated with endometriosis. A case report. *J Reprod Med* 2000;45:688-90.
86. Jeanes AC, Murray D, Davidson B, Hamilton M, Watkinson AF. Case report: hepatic and retro-peritoneal endometriosis presenting as obstructive jaundice with ascites: a case report and review of the literature. *Clin Radiol* 2002;57:226-9.
87. Francis M, Badero OO, Borowsky M, Lee YC, Abulafia O. Pericardial effusion, right-sided pleural effusion and ascites associated with stage IV endometriosis. A case report. *J Reprod Med* 2003;48:463-5.
88. Cheong EC, Lim DT. Massive ascites—an uncommon presentation of endometriosis. *Singapore Med J* 2003;44:98-100.
89. Donnez J, Jadoul P. Ascites and pelvic masses: an unusual case of endometriosis. *Fertil Steril* 2005;83:195-7.
90. Goumenou A, Matalliotakis I, Mahutte N, Koumantakis E. Endometriosis mimicking advanced ovarian cancer. *Fertil Steril* 2006;86:219.e23-5.
91. Palayekar M, Jenci J, Carlson JA Jr. Recurrent hemorrhagic ascites: a rare presentation of endometriosis. *Obstet Gynecol* 2007;110:521-2.
92. Zeppa P, Vetrani A, Cozzolino I, Palombini L. Endometrial glands in ascites secondary to endometriosis. *Diagn Cytopathol* 2004;30:131-2.
93. Lin JN, Lin HL, Huang CK, Lai CH, Chung HC, Liang SH, Lin HH. Endometriosis presenting as bloody ascites and shock. *J Emerg Med* 2008;doi:10.1016/j.jemermed.2008.03.031.
94. Bernstein JS, Perlow V, Brenner JJ. Massive ascites due to endometriosis. *Am J Dig Dis* 1961;6:1-6.
95. Wang PH, Liu WM, Fuh JL, Cheng MH, Chao HT. Comparison of surgery alone and combined surgical-medical treatment in the management of symptomatic uterine adenomyoma. *Fertil Steril* 2008;doi:10.1016/j.fertnstert.2008.07.1744.
96. Pabuccu R, Onalan G, Kaya C. GnRH agonist and antagonist protocols for stage I-II endometriosis and endometrioma in in vitro fertilization/intracytoplasmic sperm injection cycles. *Fertil Steril* 2007;88:832-9.
97. Fedele L, Bianchi S, Montefusco S, Frontino G, Carmignani L. A gonadotropin-releasing hormone agonist versus a continuous oral contraceptive pill in the treatment of bladder endometriosis. *Fertil Steril* 2008;90:183-4.
98. Ghosh SB, Mala YM, Tripathi R, Tyagi S. Coexisting genital tuberculosis and endometriosis presenting as abdominal mass in an infertile woman: a report of a rare case. *Fertil Steril* 2008;90:443.e5-6.
99. Lebovic DI, Mwenda JM, Chai DC, Mueller MD, Santi A, Fisseha S, D'Hooghe T. PPAR-gamma receptor ligand induces regression of endometrial explants in baboons: a prospective, randomized, placebo- and drug-controlled study. *Fertil Steril* 2007;88:1108-19.
100. Saito S, Murakami T, Suzuki K, Terada Y, Fukushima K, Moriya T. Intestinal endometriosis complicated by ileal perforation after initiation of gonadotropin-releasing hormone agonist therapy. *Fertil Steril* 2007;88:969.e7-9.