

FETAL MAGNETIC RESONANCE IMAGING DEMONSTRATION OF CENTRAL NERVOUS SYSTEM ABNORMALITIES AND POLYDACTYLY ASSOCIATED WITH JOUBERT SYNDROME

Chih-Ping Chen^{1,2,3,4,5,6*}, Yi-Ning Su⁷, Jon-Kway Huang⁸, Yu-Peng Liu^{8,9}, Fuu-Jen Tsai^{4,10,11},
Chun-Kuang Yang¹², Jian-Pei Huang¹, Chen-Yu Chen¹, Pei-Chen Wu¹, Wayseen Wang^{2,13}

*Departments of ¹Obstetrics and Gynecology and ²Medical Research, Mackay Memorial Hospital, Taipei,
³Department of Biotechnology, Asia University, ⁴School of Chinese Medicine, College of Chinese Medicine,
China Medical University, Taichung, ⁵Institute of Clinical and Community Health Nursing, National
Yang-Ming University, ⁶Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming
University, ⁷Department of Medical Genetics, National Taiwan University Hospital, ⁸Department of
Radiology, Mackay Memorial Hospital, ⁹Mackay Medicine, Nursing and Management College,
Taipei, Departments of ¹⁰Medical Genetics and ¹¹Medical Research, China Medical University
Hospital, Taichung, ¹²Department of Obstetrics and Gynecology, Tzu-Chi General Hospital,
Taipei Branch, and ¹³Department of Bioengineering, Tatung University, Taipei, Taiwan.*

A 33-year-old woman, gravida 2, para 1, was referred to hospital for prenatal imaging evaluation of the fetal brain and limbs at 27 weeks' gestation. The woman and her husband were healthy and non-consanguineous. Two years previously, she delivered a female baby at 26 weeks' gestation with Joubert syndrome (JBTS), Dandy-Walker malformation, ventriculomegaly, cerebellar vermis hypoplasia and polydactyly of the feet. JBTS was confirmed by magnetic resonance imaging (MRI). The infant's karyotype was 46,XX. During this pregnancy, cerebellar hypoplasia was noted at 17 weeks' gestation, and ventriculomegaly and polydactyly were noted at 18 weeks' gestation. Level II ultrasound examination at 27 weeks' gestation revealed a male fetus with a femur length and an abdominal circumference equivalent to 27 weeks, polyhydramnios with an amniotic fluid index of 24.07 cm. Macrocephaly was present with an increased biparietal distance of 7.9 cm, an increased head circumference of 28.82 cm, an increased lateral ventricle posterior horn width of 2.64 cm, an increased transcerebellar distance of 3.61 cm, an increased cisterna magna length of 2.10 cm, an increased head

circumference to femur length ratio of 5.66 (all > 95th centile). Ventriculomegaly, vermian hypoplasia, postaxial polydactyly of the left hand and preaxial polydactyly of the hallux of the left foot were also present. A tentative diagnosis of recurrent JBTS was made. Fetal MRI examination at 27 weeks' gestation revealed the "molar tooth sign", Dandy-Walker malformation, ventriculomegaly



Figure 1. A male fetus at 27 weeks' gestation with abnormal magnetic resonance imaging findings suggestive of Joubert syndrome. Prenatal axial imaging at the pontomesencephalic junction shows the "molar tooth sign" with cerebellar vermis hypoplasia, a dilated fourth ventricle (*), thickened and elongated superior cerebellar peduncles (arrows), and a deep interpeduncular fossa (arrowhead).



*Correspondence to: Dr Chih-Ping Chen, Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.
E-mail: cpc_mmh@yahoo.com
Accepted: January 27, 2010




Figure 2. Prenatal axial imaging shows a vermian cleft (arrow).



Figure 3. Prenatal axial imaging shows ventriculomegaly with dilation of the lateral ventricles.




Figure 4. Prenatal sagittal imaging shows a dilated fourth ventricle (arrowhead) with a round roof, a hypoplastic cerebellum (arrow), and a large posterior fossa cyst (*).




Figure 5. (A, B) Preaxial polydactyly (arrow) of the hallux of the foot.

and polydactyly (Figures 1–5). On the transaxial scan, there was a complex brainstem malformation consisting of cerebellar vermian hypoplasia, thickened and elongated superior cerebellar peduncles and a deep interpeduncular fossa, giving the appearance of the molar tooth sign which is pathognomonic for JBTS. The parents decided to continue the pregnancy. At 31⁺⁵ weeks' gestation, a 2,220 g live male baby was delivered with postaxial polydactyly of both hands, preaxial polydactyly of the left foot, and a 46,XY karyotype. The postnatal central nervous system findings were consistent with JBTS.

JBTS is a genetically heterogeneous disorder characterized by hypoplasia of the cerebellar vermis with a characteristic brainstem malformation and neuroradiologic molar tooth sign, intellectual disability, hypotonia, ataxia, tachypnea/apnea, and abnormal eye

movements [1]. JBTS and related disorders (JSRD) is classic JBTS with variable features such as central nervous system anomalies including occipital encephalocele and corpus callosal agenesis, retinal dystrophy, ocular coloboma, oral frenulae, tongue tumors, polydactyly, cystic renal dysplasia, nephronophthisis, and congenital hepatic fibrosis [1]. JSRD is one type from the wide spectrum of ciliopathies known to be caused by aberrant function of the primary cilia [1–4].

In the case presented here, JBTS involved an affected female and an affected male from the same sibship and is likely inherited in an autosomal recessive pattern. To date, at least nine genes and one additional locus have been identified in patients with JSRD [1]. The nine genes include *INPP5E* on 9q34.3 (JBTS1), *AHI1* on 6q23.3 (JBTS3), *NPHP1* on 2q13 (JBTS4), *MKS4/CEP290* on 12q21.32 (JBTS5), *MKS3/TMEM67* on 8q21.13–q22.1 (JBTS6), *MKS5/RPGRIPL* on 16q12.2 (JBTS7), *ARL13B* on 3q11.2 (JBTS8), *CC2D2A* on 4p15.3

(JBTS9), and *CXORF5* on Xp22.3 (JBTS10). The additional locus of *CORS2* has been suggested on 11p12-q13.3 (JBTS2). JBTS1–9 are inherited in an autosomal recessive pattern. JBTS10 is inherited in an X-linked recessive pattern [5]. Meckel syndrome (MKS) shares JBTS phenotypes [6,7]. Among the nine JSRD genes, three genes (*MKS3/TMEM67*, *MKS4/CEP290* and *MKS5/RPGRIPL1*) are also associated with MKS, indicating genetic complexity in JSRD. Both of the affected siblings in this case had polydactyly. Polydactyly is present in 8% [8] to 15% [9] of subjects with JSRD. Polydactyly has been shown to be present in cases with JBTS2, JBTS6 and JBTS7 but has rarely been described in other types of JSRD [1]. Polydactyly associated with JSRD is often postaxial, although preaxial polydactyly of the hands or large toes has been observed [1].

JSRD has an estimated prevalence of 1 in 100,000 live births [1]. Prenatal imaging of JSRD has rarely been described in the literature. To date, only 14 reports of prenatal imaging of JSRD have been described [10–23], most of which were examined by ultrasound and only three reports [21–23] were examined by fetal MRI. Prenatal sonographic findings in fetuses with JSRD are relatively nonspecific and include increased nuchal translucency, enlarged cisterna magna, cerebellar vermian agenesis, occipital encephalocele, ventriculomegaly, hypoplastic phallus, renal cysts, and polydactyly [20,21]. Prenatal ultrasound lacks sensitivity with regard to posterior fossa malformations. In contrast, fetal MRI is able to identify the characteristic molar tooth sign, which is the cardinal diagnostic imaging sign of JSRD. The molar tooth sign consists of cerebellar vermian hypoplasia, thickened and elongated superior cerebellar peduncles and a deep interpeduncular fossa, and mid-brain dysgenesis is responsible for the molar tooth sign [24–26]. Fluss et al [22] identified the molar tooth sign at 27 weeks of gestation by fetal MRI in a fetus with JBTS. Saleem and Zaki [23] were able to identify the molar tooth sign associated with JSRD as early as 22 weeks of gestation by fetal MRI. We suggest that fetal MRI should be considered in pregnancy with a previous JSRD-affected child and/or documented posterior fossa malformations with associated anomalies suspicious of JSRD.

Acknowledgments

This work was supported by research grants NSC-96-2314-B-195-008-MY3 and NSC-97-2314-B-195-006-MY3 from the National Science Council, and MMH-E-99004 from Mackay Memorial Hospital, Taipei, Taiwan.

References

1. Parisi MA. Clinical and molecular features of Joubert syndrome and related disorders. *Am J Med Genet C Semin Med Genet* 2009;151C:326–40.
2. Adams M, Smith UM, Logan CV, Johnson CA. Recent advances in the molecular pathology, cell biology and genetics of ciliopathies. *J Med Genet* 2008;45:257–67.
3. Baker K, Beales PL. Making sense of cilia in disease: the human ciliopathies. *Am J Med Genet C Semin Med Genet* 2009;151C:281–95.
4. Cardenas-Rodriguez M, Badano JL. Ciliary biology: understanding the cellular and genetic basis of human ciliopathies. *Am J Med Genet C Semin Med Genet* 2009;151C:263–80.
5. Coene KLM, Roepman R, Doherty D, et al. OFD1 is mutated in X-linked Joubert syndrome and interacts with LCAS-encoded lebercilin. *Am J Hum Genet* 2009;85:465–81.
6. Chen CP. Meckel syndrome: genetics, perinatal findings, and differential diagnosis. *Taiwan J Obstet Gynecol* 2007;46:9–14.
7. Chen CP. Syndromes, disorders and maternal risk factors associated with neural tube defects (III). *Taiwan J Obstet Gynecol* 2008;47:131–40.
8. Chance PF, Cavalier L, Satran D, Pellegrino JE, Koenig M, Dobyns WB. Clinical nosologic and genetic aspects of Joubert and related syndromes. *J Child Neurol* 1999;14:660–6.
9. Saraiva JM, Baraitser M. Joubert syndrome: a review. *Am J Med Genet* 1992;43:726–31.
10. Campbell S, Tsannatos C, Pearce JM. The prenatal diagnosis of Joubert's syndrome of familial agenesis of the cerebellar vermis. *Prenat Diagn* 1984;4:391–5.
11. van Drop DB, Palan A, Kwee ML, Barth PG, van der Harten JJ. Joubert syndrome: clinical and pathological description of an affected male and a female fetus from the sibship. *Am J Med Genet* 1991;40:100–4.
12. Ivarsson SA, Bjerre I, Brun A, Ljungberg O, Maly E, Taylor I. Joubert syndrome associated with Leber amaurosis and multicystic kidneys. *Am J Med Genet* 1993;45:542–7.
13. Van Zalen-Sprock RM, van Vugt JM, van Geijn HP. First-trimester sonographic detection of neurodevelopmental abnormalities in some single-gene disorders. *Prenat Diagn* 1996;16:199–202.
14. Reynders CS, Pauker SP, Benacerraf BR. First trimester isolated fetal nuchal lucency: significance and outcome. *J Ultrasound Med* 1997;16:101–5.
15. Souka AP, Snijders RJM, Novakov A, Soares W, Nicolaides KH. Defects and syndromes in chromosomally normal fetuses with increased nuchal translucency thickness at 10–14 weeks of gestation. *Ultrasound Obstet Gynecol* 1998;11:391–400.
16. Anderson JS, Gorey MT, Pasternak JF, Trommer BL. Joubert's syndrome and prenatal hydrocephalus. *Pediatr Neurol* 1999;20:403–5.
17. Ni Scanail S, Crowley P, Hogan M, Stuart B. Abnormal prenatal sonographic findings in the posterior cranial fossa: a case of Joubert's syndrome. *Ultrasound Obstet Gynecol* 1999;13:71–4.
18. Wang P, Chang FM, Chang CH, Yu CH, Jung YC, Huang CC. Prenatal diagnosis of Joubert syndrome complicated with encephalocele using two-dimensional and three-dimensional ultrasound. *Ultrasound Obstet Gynecol* 1999;14:360–2.

19. Aslan H, Ulker V, Gulcan EM, Numanoglu C, Gul A, Agar M, Ark HC. Prenatal diagnosis of Joubert syndrome: a case report. *Prenat Diagn* 2002;22:13–6.
20. Aslan H, Gungorduk K, Yildirim G, Olgac Y, Ceylan Y. Prenatal ultrasonographic features of Joubert syndrome. *J Clin Ultrasound* 2008;36:576–80.
21. Doherty D, Glass IA, Siebert JR, et al. Prenatal diagnosis in pregnancies at risk for Joubert syndrome by ultrasound and MRI. *Prenat Diagn* 2005;25:442–7.
22. Fluss J, Blaser S, Chitayat D, Akoury H, Glanc P, Skidmore M, Raybaud C. Molar tooth sign in fetal brain magnetic resonance imaging leading to the prenatal diagnosis of Joubert syndrome and related disorders. *J Child Neurol* 2006;21:320–4.
23. Saleem SN, Zaki MS. Role of MR imaging in prenatal diagnosis of pregnancies at risk for Joubert syndrome and related cerebellar disorders. *Am J Neuroradiol* 2010;31:424–9.
24. Maria BL, Hoang KB, Tusa RJ, et al. “Joubert syndrome” revisited: key ocular motor signs with magnetic resonance imaging correlation. *J Child Neurol* 1997;12:423–30.
25. Maria BL, Quisling RG, Rosainz LC, Yachnis AT, Gitten J, Dede D, Fennell E. Molar tooth sign in Joubert syndrome: clinical, radiologic, and pathologic significance. *J Child Neurol* 1999;14:368–76.
26. Maria BL, Boltshauser E, Palmer SC, Tran TX. Clinical features and revised diagnostic criteria in Joubert syndrome. *J Child Neurol* 1999;14:583–90.