

APERT SYNDROME ASSOCIATED WITH UPPER AIRWAY OBSTRUCTION AND GASTROESOPHAGEAL REFLUX INDUCING POLYHYDRAMNIOS IN THE THIRD TRIMESTER

Chih-Ping Chen^{1,2,3,4,5,6*}, Shuan-Pei Lin^{2,7}, Yi-Ning Su⁸, Chen-Yu Chen¹, Fuu-Jen Tsai^{4,9,10}, Yu-Peng Liu^{11,12}, Schu-Rern Chern², Pei-Chen Wu¹, Hsiao-En Cindy Chen¹³, Wayseen Wang^{2,14}

Departments of ¹Obstetrics and Gynecology and ²Medical Research, Mackay Memorial Hospital, Taipei, ³Department of Biotechnology, Asia University, ⁴School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, ⁵Institute of Clinical and Community Health Nursing, National Yang-Ming University, ⁶Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, ⁷Department of Pediatrics, Mackay Memorial Hospital, ⁸Department of Medical Genetics, National Taiwan University Hospital, Taipei, Departments of ⁹Medical Genetics and ¹⁰Medical Research, China Medical University Hospital, Taichung, ¹¹Department of Radiology, Mackay Memorial Hospital, ¹²Mackay Medicine, Nursing and Management College, Taipei, Taiwan, ¹³School of Medicine, University of Wollongong, New South Wales, Australia, and ¹⁴Department of Bioengineering, Tatung University, Taipei, Taiwan.

Here, we present the case of a female infant, the first child of a 32-year-old woman and a 45-year-old man, who were healthy and non-consanguineous. The family history was unremarkable. The infant was delivered at 37 weeks' gestation with a birth weight of 2,834 g. She had midface hypoplasia, cleft palate, low-set ears, and bilateral syndactyly of the hands and feet (Figures 1–3). She had a 46,XX karyotype. DNA testing for Apert syndrome revealed a heterozygous c.755 C > G, TCG > TGG transversion leading to a Ser252Trp (S252W) mutation in the fibroblast growth factor receptor 2 (*FGFR2*; OMIM 176943) gene. The pregnancy was uneventful until the third trimester when polyhydramnios was noted (Figure 4). The fetal brain, skull shape and internal organs were normal. The amniotic fluid index was measured as 37.4 cm, 32.3 cm and 30.4 cm at 33, 35 and 36 weeks of gestation, respectively. After birth, the infant manifested respiratory distress owing to a decreased nasopharyngeal airway. Echocardiography showed an atrial septal defect. Three-dimensional computed tomography (CT) reconstruction of the skull showed brachycephaly, shallow bilateral orbital fossa due to premature synostosis of the bilateral coronal sutures, and

midface hypoplasia (Figure 5). CT showed normal brain parenchyma, a normal ventricular system, maxillary hypoplasia, symmetric narrowing of both nasal cavities, choanal stenosis, and a narrow and high-arched palate (Figures 6 and 7). The infant underwent tracheostomy at 2 months of age for the treatment of cyanotic episodes, obstructive stridor, and sleep apnea. Bronchoscopy performed prior to surgery revealed upper airway obstruction with nasal stenosis, tongue drop, and laryngomalacia. The subglottis, glottis, trachea and bronchus were normal. She underwent gastrostomy, pyloroplasty and fundoplication at 12 months of age for the treatment of gastroesophageal reflux, frequent vomiting, and poor feeding. She underwent first-stage hand surgery at 2 years of age and second-stage hand surgery at 3 years of age for the treatment of bilateral syndactyly.

Apert syndrome (OMIM 101200) is characterized by acrocephaly, craniosynostosis, a flat occiput, midface hypoplasia, and "mitten-like" syndactyly of the hands and feet [1]. The prevalence at birth of Apert syndrome is estimated to be 1 in 160,000 to 1 in 164,500 [2,3]. Two mutations of the *FGFR2* gene, Ser252Trp (S252W) and Pro253Arg (P253R), account for over 98% of cases with Apert syndrome [4,5]. Most cases of Apert syndrome arise *de novo* but autosomal dominant inheritance and germline mosaicism have been described [6]. Advanced paternal age can be a risk factor [2,7,8]. Although cases of prenatally diagnosed Apert syndrome have been reported [9], prenatal diagnosis of Apert syndrome remains difficult. Many cases of Apert



ELSEVIER

* Correspondence to: Dr Chih-Ping Chen, Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.

E-mail: cpc_mmh@yahoo.com

Accepted: January 12, 2010



Figure 1. “Mitten-like” syndactyly of the hands.



Figure 2. “Mitten-like” syndactyly of the feet.



Figure 3. “Mitten-like” syndactyly of the feet.

syndrome were undiagnosed until delivery, or diagnosed in late gestation when polyhydramnios and craniofacial deformities became evident.

The present case had the S252W mutation. von Gernet et al [10] observed more severe midface and dental findings in S252W patients than in P253R patients, and conversely, more severe syndactyly of hands in P253R patients than in S252W patients. Airway obstruction has been observed in 40% of patients with severe craniosynostotic syndromes [11,12]. In patients with Apert syndrome and midface hypoplasia, the maxilla is narrow and retruded, resulting in a narrow and high-arched palate, small postnasal space and




Figure 4. (A) Lateral and (B) anterior views on prenatal ultrasound showing a depressed nasal bridge and midface hypoplasia.




Figure 5. Three-dimensional computed tomography reconstruction of the skull.

clivus, narrowing of the bony nasal cavities, and choanal stenosis [13]. Airway obstruction may be secondary to midface hypoplasia and may result in sleep apnea and cor pulmonale that require tracheostomy [11,14–18].

Polyhydramnios may develop in the third trimester in pregnancies with fetal Apert syndrome [19]. Nyberg et al [20] suggested that polyhydramnios associated with Apert syndrome is developed from decreased fetal swallowing related to central nervous system abnormalities. However, the present case did not have central nervous system abnormalities. The present case had upper airway obstruction which can cause decreased fetal swallowing. The present case provides evidence that a reduced nasopharyngeal airway can be a principal causative factor of polyhydramnios in pregnancies with fetal Apert syndrome. Fetal Apert syndrome associated with polyhydramnios in the third trimester should be indicative of upper airway obstruction. In such cases, prompt early diagnosis and intervention after birth are necessary to reduce perinatal morbidity.




Figure 6. (A) Axial and (B) coronal computed tomography scans of the nose and the paranasal sinuses showing symmetric narrowing of both nasal cavities.




Figure 7. Sagittal computed tomography scan of the skull showing brachycephaly, midface hypoplasia, a narrow and retruded maxilla, a high-arched palate, and a decreased nasopharyngeal airway.

Acknowledgments

This work was supported by research grants NSC-96-2314-B-195-008-MY3 and NSC-97-2314-B-195-006-MY3 from the National Science Council, and MMH-E-98004 from Mackay Memorial Hospital, Taipei, Taiwan.

References

1. Apert ME. De l'acrocephalosyndactylie. *Bull Mem Soc Med Hop Paris* 1906;23:1310–30. [In French]
2. Blank CE. Apert's syndrome (a type of acrocephalosyndactyly): observations on a British series of thirty-nine cases. *Ann Hum Genet* 1960;24:151–64.
3. Cohen MM Jr, Kreiborg S, Lammer EJ, et al. Birth prevalence study of the Apert syndrome. *Am J Med Genet* 1992;42:655–9.
4. Park WJ, Theda C, Maestri NE, et al. Analysis of phenotypic features and FGFR2 mutations in Apert syndrome. *Am J Hum Genet* 1995;57:321–8.
5. Wilkie AO, Slaney SF, Oldridge M, et al. Apert syndrome results from localized mutations of FGFR2 and is allelic with Crouzon syndrome. *Nat Genet* 1995;9:165–72.
6. Allanson JE. Germinal mosaicism in Apert syndrome. *Clin Genet* 1986;29:429–33.
7. Erickson JD, Cohen MM Jr. A study of parental age effects on the occurrence of fresh mutations for the Apert syndrome. *Ann Hum Genet* 1974;38:89–96.
8. Moloney DM, Slaney SF, Oldridge M, Wall SA, Sahlin P, Stenman G, Wilkie AO. Exclusive paternal origin of new mutations in Apert syndrome. *Nat Genet* 1996;13:48–53.
9. Chen CP, Su YN, Hsu CY, et al. Second-trimester molecular prenatal diagnosis of sporadic Apert syndrome following sonographic findings of mild ventriculomegaly and clenched hands mimicking trisomy 18. *Taiwan J Obstet Gynecol* 2010;49:129–32.
10. von Gernet S, Golla A, Ehrenfels Y, Schuffenhauer S, Fairley JD. Genotype-phenotype analysis in Apert syndrome suggests opposite effects of the two recurrent mutations on syndactyly and outcome of craniofacial surgery. *Clin Genet* 2000;57:137–9.
11. Sculerati N, Gottlieb MD, Zimble MS, Chibbaro PD, McCarthy JG. Airway management in children with major craniofacial anomalies. *Laryngoscope* 1998;108:1806–12.
12. Lo LJ, Chen YR. Airway obstruction in severe syndromic craniosynostosis. *Ann Plast Surg* 1999;43:258–64.
13. McGill T. Otolaryngologic aspects of Apert syndrome. *Clin Plast Surg* 1991;18:309–13.
14. Schafer ME. Upper airway obstruction and sleep disorders in children with craniofacial anomalies. *Clin Plast Surg* 1982;9:555–67.
15. Lauritzen C, Lilja J, Jarlstedt J. Airway obstruction and sleep apnea in children with craniofacial anomalies. *Plast Reconstr Surg* 1986;77:1–6.

16. Kaplan LC. Clinical assessment and multispecialty management of Apert syndrome. *Clin Plast Surg* 1991;18:217–25.
17. Järund M, Lauritzen C. Craniofacial dysostosis: airway obstruction and craniofacial surgery. *Scand J Plast Reconstr Surg Hand Surg* 1996;30:275–9.
18. Perkins JA, Sie KC, Milczuk H, Richardson MA. Airway management in children with craniofacial anomalies. *Cleft Palate Craniofac J* 1997;34:135–40.
19. Filkins K, Russo JF, Boehmer S, Camous M, Przylepa KA, Jiang W, Jabs EW. Prenatal ultrasonographic and molecular diagnosis of Apert syndrome. *Prenat Diagn* 1997;17:1081–4.
20. Nyberg DA, Jeanty P, Glass I. Syndromes and multiple anomaly conditions. In: Nyberg DA, McGahan JP, Pretorius DH, Pilu G, eds. *Diagnostic Imaging of Fetal Anomalies*. Philadelphia: Lippincott Williams & Wilkins, 2003:133–220.