

Research Letter

# Prenatal diagnosis and molecular cytogenetic characterization of a proximal deletion of 22q (22q11.2 → q11.21)

Chih-Ping Chen<sup>a,b,c,d,e,f,g,\*</sup>, Tsang-Ming Ko<sup>h</sup>, Yi-Ning Su<sup>i</sup>, Jun-Wei Su<sup>b,j</sup>, Yu-Ting Chen<sup>c</sup>,  
Chen-Chi Lee<sup>b</sup>, Li-Feng Chen<sup>b</sup>, Wayseen Wang<sup>c,k</sup>

<sup>a</sup> Department of Medicine, Mackay Medical College, New Taipei City, Taiwan

<sup>b</sup> Department of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, Taiwan

<sup>c</sup> Department of Medical Research, Mackay Memorial Hospital, Taipei, Taiwan

<sup>d</sup> Department of Biotechnology, Asia University, Taichung, Taiwan

<sup>e</sup> School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

<sup>f</sup> Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan

<sup>g</sup> Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

<sup>h</sup> GenePhile Bioscience Laboratory, Ko's Obstetrics and Gynecology, Taipei, Taiwan

<sup>i</sup> Department of Obstetrics and Gynecology, School of Medicine, College of Medicine, Taipei Medical University, Taipei, Taiwan

<sup>j</sup> Department of Obstetrics and Gynecology, China Medical University Hospital, Taichung, Taiwan

<sup>k</sup> Department of Bioengineering, Tatung University, Taipei, Taiwan

Accepted 29 September 2012

A gravid 2, para 1, woman 34 years of age underwent amniocentesis at 18 weeks of gestation because of advanced maternal age. She had a healthy daughter 3 years of age, and there was no family history of congenital malformations. Amniocentesis revealed a karyotype of 45,XX,der(13)t(13;22)(q10;q11.2),-22 (Fig. 1). The father had a karyotype of 46,XY, the mother had a karyotype of 46,XX,t(13;22)(q10;q11.2) (Fig. 2), and the daughter had a karyotype of 46,XX. Prenatal ultrasonographic findings of the internal organs, including the heart, thymus, brain, and kidneys, were unremarkable. Oligonucleotide-based array comparative genomic hybridization (aCGH) analysis of cultured amniocytes using CytoChip Oligo array (BlueGnome, Cambridge, UK) revealed a 1.59-Mb deletion at chromosome 22q11.1-22q11.21 or arr 22q11.1q11.21 (17,397,528–18,984,490 bp) × 1 (National Center for Biotechnology Information (NCBI), Build 37; Fig. 3). The deleted region of proximal 22q was proximal to the DiGeorge syndrome (DGS) critical region. Fluorescence *in situ* hybridization (FISH) analysis of cultured amniocytes using Vysis DiGeorge region probe [Vysis, LSI TUPLE 1 (red spectrum)/LSI ARSA (green spectrum)] (Abbott Laboratories, Abbott Park, IL, USA) showed the presence of two red and two green signals, indicating no deletion of the DGS TUPLE 1

locus at 22q11.2 (Fig. 4). The woman underwent cord blood sampling at 24 weeks of gestation. The cord blood was analyzed by multiplex ligation-dependent probe amplification (MLPA) using the SALSA MLPA P250 DiGeorge probemix (MRC-Holland bv. Amsterdam, The Netherlands) to detect cat eye syndrome (22q11; CES), DiGeorge syndrome (22q11.2, LCR22-A~G), velocardiofacial syndrome (22q11; VCFS) and the deletions of DiGeorge anomaly related regions such as 4q35, 8p23, 9q34.3, 10p14(DGS2), and 17p13. The SALSA MLPA P250 DiGeorge probemix showed haploinsufficiency of genes *IL17RA*, *SLC25A18*, *BID*, *MICAL3*, and *USP18* in the CES region but no deletion in the DGS critical region of *TBX1* or any other DiGeorge anomaly-related regions (Fig. 5). The parents decided to continue the pregnancy. At 39 weeks of gestation, a healthy 3,474-g female baby was delivered with no phenotypic abnormalities. At 3 months of age, the neonate was normal in growth and psychomotor development although long-term follow-up visits are required.

MLPA is a useful tool of rapid aneuploidy diagnosis without the need of cell culture. We previously reported the applications of the SALSA MLPA P095 aneuploidy kit for rapid aneuploidy diagnosis of chromosomes X, Y, 13, 18, and 21 in pregnancies with fetal congenital anomalies [1,2]. Here, we demonstrate the usefulness of the SALSA MLPA P250 DiGeorge probemix as well as FISH and aCGH for the diagnosis of proximal 22q deletion without involvement of the DGS critical region in a case with an unbalanced t(13;22) translocation involving chromosome 22q11.2. MLPA has

\* Corresponding author. Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.

E-mail address: [cpc\\_mmh@yahoo.com](mailto:cpc_mmh@yahoo.com) (C.-P. Chen).




Fig. 1. A karyotype of 45,XX,der(13)t(13;22)(q10;q11.2) in the fetus.

proven to be a rapid, reliable, and sensitive tool for detecting copy number changes in the 22q11.2 region, including DGS, VCFS, conotruncal anomaly face syndrome, and CES [3–5]. The present case had a 1.59-Mb deletion at chromosome 22q11.1–22q11.21 encompassing the CES critical region genes of *CECR1* (OMIM 607575) (17,660,191–17,690,778 bp) and *CECR2* (OMIM 607576) (17,956,629–18,033,844 bp) but without involvement of the DGS critical region gene of *TBX1* (OMIM 602054) (19,744,225–19,771,115 bp) at 22q11.21.

CES (OMIM 115470) is characterized by coloboma of iris, anal atresia with fistula, preauricular tags and/or pits, down-slanting palpebral fissures, cardiac and renal malformations, and normal or near-normal mental development. CES is frequently associated with partial tetrasomy 22q or a supernumerary chromosome of inv dup(22)(q11). Deletions of the juxtacentromeric region or chromosome 22 have been reported to be associated with phenotypic variability [6], ranging from phenotypically normal [7,8] to mild and relatively nonspecific




Fig. 2. A karyotype of 46,XX,t(13;22)(q10;q11.2) in the mother.




Fig. 3. Oligonucleotide-based array comparative genomic hybridization analysis of cultured amniocytes using CytoChip Oligo array (BlueGnome, Cambridge, UK) revealed a 1.59-Mb deletion at chromosome 22q11.1-22q11.21 or arr 22q11.1q11.21 (17,397,528–18,984,490 bp)  $\times$ 1 (NCBI, Build 37).




Fig. 4. (A) Fluorescence *in situ* hybridization analysis using the Vysis DiGeorge region probe [Vysis, LSI TUPLE 1 (red spectrum)/LSI ARSA (green spectrum)] (Abbott Laboratories, Abbott Park, IL, USA) showed the presence of two red signals and two green signals in interphase cultured amniocytes; and (B) metaphase cultured amniocytes, indicating no deletion of the DiGeorge syndrome TUPLE 1 locus at 22q11.2 in the fetus. der = derivative chromosome.



## References

- [1] Chen C-P, Su Y-N, Lin S-Y, Chang C-L, Wang Y-L, Huang J-P, et al. Rapid aneuploidy diagnosis by multiplex ligation-dependent probe amplification and array comparative genomic hybridization in pregnancy with major congenital malformations. *Taiwan J Obstet Gynecol* 2011;50: 85–94.
- [2] Chen C-P, Su Y-N, Chang C-L, Chen Y-Y, Su J-W, Chern S-R, et al. Rapid aneuploidy diagnosis by multiplex ligation-dependent probe amplification using uncultured amniocytes in pregnancy with major fetal structural abnormalities. *Taiwan J Obstet Gynecol* 2012;51:123–8.
- [3] Fernández L, Lapunzina P, Arjona D, López Pajares I, García-Guereta L, Elorza D, et al. Comparative study of three diagnostic approaches (FISH, STRs and MLPA) in 30 patients with 22q11.2 deletion syndrome. *Clin Genet* 2005;68:373–8.
- [4] Vorstman JAS, Jalali GR, Rappaport EF, Hacker AM, Scott C, Emanuel BS. MLPA: a rapid, reliable, and sensitive method for detection and analysis of abnormalities of 22q. *Hum Mutat* 2006;27:814–21.
- [5] Jalali GR, Vorstman JA, Errami A, Vijzelaar R, Biegel J, Shaikh T, et al. Detailed analysis of 22q11.2 with a high density MLPA probe set. *Hum Mutat* 2008;29:433–40.
- [6] Schinzel A. del(22)(pter→q11.2). In: Schinzel A, editor. Catalogue of unbalanced chromosome aberrations in man. 2nd ed. Berlin: Walter de Gruyter, GmbH & Co.; 2001. p. 843–5.
- [7] Kirkels VGHJ, Hustinx TWJ, Scheres JMJC. Habitual abortion and translocation (22q;22q): unexpected transmission from a mother to her phenotypically normal daughter. *Clin Genet* 1980;18:456–61.
- [8] Palmer CG, Schwartz S, Hodes ME. Transmission of a balanced homologous t(22q;22q) translocation from mother to normal daughter. *Clin Genet* 1980;17:418–22.
- [9] Nakagome Y, Inuma K, Matsui I. Three translocations involving C- or G-group chromosomes. *J Med Genet* 1973;10:174–6.
- [10] Jenkins MB, Davis E, Thelen TH, Boyd L. A familial X-22 translocation with an extra X chromosome. *Am J Hum Genet* 1974;26:736–45.
- [11] Abe T, Morita M, Misawa S, Hattori A, Nakano S, Hojo H. Breakage and reunion at the chromosome band 22q11 in somatic cells — its relevance to the genetic role of the Philadelphia chromosome. *Nihon Ketsueki Gakkai Zasshi* 1978;41:140–3.
- [12] Vitelli F, Viola A, Morishima M, Pramparo T, Baldini A, Lindsay E. *TBX1* is required for inner ear morphogenesis. *Hum Mol Genet* 2003;12: 2041–8.
- [13] Lichtner P, König R, Hasegawa T, Van Esch H, Meitinger T, Schuffenhauer S. An HDR (hypoparathyroidism, deafness, renal dysplasia) syndrome locus maps distal to the DiGeorge syndrome region on 10p13/14. *J Med Genet* 2000;37:33–7.
- [14] Villanueva MP, Aiyer AR, Muller S, Pletcher MT, Liu X, Emanuel B, et al. Genetic and comparative mapping of genes dysregulated in mouse hearts lacking the *Hand2* transcription factor gene. *Genomics* 2002;80: 593–600.
- [15] Yatsenko SA, Yatsenko AN, Szigeti K, Craigen WJ, Stankiewicz P, Cheung SW, et al. Interstitial deletion of 10p and atrial septal defect in DiGeorge 2 syndrome. *Clin Genet* 2004;66:128–36.
- [16] Devriendt K, Matthijs G, Van Dael R, Gewillig M, Eyskens B, Hjalgrim H, et al. Delineation of the critical deletion region for congenital heart defects, on chromosome 8p23.1. *Am J Hum Genet* 1999;64: 1119–26.
- [17] Tsai C-H, Van Dyke DL, Feldman GL. Child with velocardiofacial syndrome and del(4)(q34.2): another critical region associated with a velocardiofacial syndrome-like phenotype. *Am J Med Genet* 1999;82: 336–9.
- [18] Kleefstra T, van Zelst-Stams WA, Nillesen WM, Cormier-Daire V, Houge G, Foulds N, et al. Further clinical and molecular delineation of the 9q subtelomeric deletion syndrome supports a major contribution of *EHMT1* haploinsufficiency to the core phenotype. *J Med Genet* 2009;46: 598–606.
- [19] Greenberg F, Courtney KB, Wessels RA, Huhta J, Carpenter RJ, Rich DC, et al. Prenatal diagnosis of deletion 17p13 associated with DiGeorge anomaly. *Am J Med Genet* 1988;31:1–4.