

Case Report

Prenatal diagnosis and molecular cytogenetic characterization of a *de novo* interstitial duplication of 14q (14q31.3 → q32.12) associated with abnormal maternal serum biochemistry

Chih-Ping Chen^{a,b,c,d,e,f,g,*}, Kwui-Shuai Hwang^h, Her-Young Su^h, Shuan-Pei Lin^{b,c,i,j},
Yi-Ning Su^k, Schu-Rern Chern^b, Jun-Wei Su^{a,l}, Yu-Ting Chen^b, Wen-Lin Chen^a,
Wayseen Wang^{b,m}

^a Department of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, Taiwan

^b Department of Medical Research, Mackay Memorial Hospital, Taipei, Taiwan

^c Department of Medicine, Mackay Medical College, New Taipei City, Taiwan

^d Department of Biotechnology, Asia University, Taichung, Taiwan

^e School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^f Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan

^g Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^h Department of Obstetrics and Gynecology, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan

ⁱ Department of Pediatrics, Mackay Memorial Hospital, Taipei, Taiwan

^j Mackay Medicine, Nursing and Management College, Taipei, Taiwan

^k Department of Obstetrics and Gynecology, School of Medicine, College of Medicine, Taipei Medical University, Taipei, Taiwan

^l Department of Obstetrics and Gynecology, China Medical University Hospital, Taichung, Taiwan

^m Department of Bioengineering, Tatung University, Taipei, Taiwan

Accepted 27 August 2012

Abstract

Objective: To present prenatal diagnosis and molecular cytogenetic characterization of a *de novo* interstitial duplication of 14q (14q31.3 → q32.12) in a pregnancy associated with abnormal maternal serum biochemistry.

Case Report: A 19-year-old woman underwent amniocentesis in the second trimester because of abnormal maternal serum biochemistry. Her husband was 33 years old. At 16 weeks of gestation, the levels of α -fetoprotein, unconjugated estriol, total β -human chorionic gonadotropin, and inhibin A were 0.8 multiples of median (MoM), 0.84 MoM, 3.06 MoM, and 1.14 MoM, respectively, consistent with a positive trisomy 21 risk of 1/269. Results of an amniocentesis revealed a small *de novo* interstitial duplication of 14q encompassing 14q31-q32.1. An array comparative genomic hybridization analysis detected a 6.6-Mb duplication at chromosome 14q31.3-q32.12. Results of a fluorescence *in situ* hybridization analysis showed a direct duplication of interstitial 14q. The karyotype was 46,XY,dup(14)(q31.3q32.12). Level II ultrasound was unremarkable. The parents decided to continue the pregnancy. A 3805-g healthy male baby was delivered at 39 weeks of gestation. When examined at 6 months of age, the neonate was normal in growth and psychomotor development with no apparent phenotypic abnormalities, although long-term follow-ups are required.

Conclusion: Abnormal maternal serum biochemistry in the second trimester may be a distinctive prenatal feature in pregnancy associated with fetal chromosome 14q duplication.

Copyright © 2013, Taiwan Association of Obstetrics & Gynecology. Published by Elsevier Taiwan LLC. All rights reserved.

Keywords: 14q31.3 → q32.12; duplication 14q; interstitial duplication; maternal serum biochemistry; prenatal diagnosis

* Corresponding author. Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.
E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).

Introduction

Partial trisomy 14q with a duplication of 14q (14q22→q32) has been associated with a distinct 14q duplication syndrome characterized by craciofacial dysmorphisms of high forehead, wide sutures and fontanels, sparse hair and eyebrows, broad nasal bridge, hypertelorism, thin upper lip, abnormal ears, broad mouth and micrognathia, developmental delay, mild mental retardation, spasticity, hyperreflexia, premature puberty, and primordial short stature [1–5].

Cases of partial trisomy 14q with a duplication of 14q31→qter are rare and present only minor anomalies of an asymmetric face, frontal bossing, low-set and dysplastic ears, hypertelorism, sparse eyebrows and lashes, prominent nasal bridge, small mandible, growth retardation, and moderate mental retardation [6–13]. Herein, we present prenatal diagnosis and molecular cytogenetic characterization of a *de novo* interstitial duplication of 14q (14q31.3→q32.12) in a pregnancy associated with abnormal maternal serum biochemistry but without apparent phenotypic abnormalities. To our knowledge, such a case has not previously been described.

Case report

A 19-year-old, gravida 2, para 1, woman underwent second-trimester screening for chromosome abnormalities using maternal serum biochemistry at 16 weeks of gestation. Her husband was 33 years old. The levels of α -fetoprotein (AFP), unconjugated estriol (uE3), total β -human chorionic gonadotropin (β -hCG), and inhibin-A were 0.8 multiples of median (MoM), 0.84 MoM, 3.06 MoM, and 1.14 MoM, respectively, consistent with a positive trisomy 21 risk of 1/269. At 18 weeks of gestation, she underwent amniocentesis, which revealed a small interstitial duplication of 14q encompassing 14q31-q32.1 (Fig. 1). Prenatal ultrasound findings were unremarkable. A repeated amniocentesis was performed at 20 weeks of gestation.

Oligonucleotide-based array comparative genomic hybridization using CytoChip Oligo Array (BlueGnome, Cambridge, UK) on uncultured amniocytes detected a 6.6-Mb duplication at chromosome 14q31.3-q32.12, or arr cgh 14q31.3q32.12 (84,238,307–90,839,109) \times 3 (UCSC hg18, NCBI build 36, March 2006) (Fig. 2). The parental karyotypes were normal. For fluorescence *in situ* hybridization (FISH) determination of the orientation of the duplication of the chromosome 14 [dup(14)], the bacterial artificial chromosome (BAC) clone probes mapping the genomic region of 14q31.3-q32.1 were used. The BAC clone probes RP11-35P13 (85,596,133–85,731,472) (UCSC hg18, NCBI build 36) (green spectrum) at 14q31.3 and RP11-99C24 (89,927,464–90,094,722) (red spectrum) at 14q32.11 were used to determine the orientation of the duplication. A FISH analysis showed an orientation of green–red–green–red (Fig. 3), consistent with a direct duplication of interstitial 14q. Conventional cytogenetic analysis of cultured amniocytes revealed a karyotype of 46,XY,dup(14)(q31.3q32.12) (Fig. 1). Level II ultrasound was unremarkable. The parents opted to continue the pregnancy. At 39 weeks of gestation, a healthy male baby was delivered with a body weight of 3805 g (85th centile), a body length of 51 cm (85th centile), and a head circumference of 36 cm (85th centile). There was no dysmorphism. Ultrasound examinations of internal organs showed normal findings. When examined at 6 months of age, he was normal in growth and psychomotor development with no apparent phenotypic abnormalities, although long-term follow-ups are required.

Discussion

Abnormal maternal serum biochemistry tests in the first or second trimester may result in incidental detection of rare fetal chromosomal abnormalities [14–16]. The present pregnant woman was 19 years old. She underwent amniocentesis because of a positive screen risk of 1/269 for trisomy 21 calculated by relatively low levels of AFP and uE3, and an




Fig. 1. A karyotype of 46,XY,dup(14)(q31.3q32.12).




Fig. 2. Oligonucleotide-based array comparative genomic hybridization on uncultured amniocytes shows a 6.6-Mb duplication at 14q31.3-q32.12 (arrow).

abnormally high level of β -hCG. The present case shows that fetuses with an interstitial duplication of 14q may present abnormal maternal serum biochemistry in the second trimester, and suggests that abnormal maternal serum biochemistry may be a distinctive prenatal feature in pregnancy associated with fetal chromosome 14q duplication.

Our case is the first report of a *de novo* tandem duplication encompassing the chromosomal bands from 14q31.3 to 14q32.12 without involving 14q32.2. The present case did not have the phenotype of uniparental disomy (UPD) 14 and 14q duplication syndrome. Maternal UPD 14 manifests a phenotype of short stature, growth retardation, muscular hypotonia,

joint laxity, truncal obesity, small hands, hyperextensible joints, scoliosis, and mild dysmorphic features of the face, whereas paternal UPD 14 manifests intrauterine growth restriction, polyhydramnios, severe psychomotor retardation, mild contractures of the fingers, cardiomyopathy, and the “coat-hanger sign” of the thoracic ribs [17,18]. Genomic imprinting effects have been observed in chromosome 14 because the 14q32.2 region in human chromosome harbors paternally expressed genes such as *DLK1* and *RTL1*, maternally expressed genes such as *MEG3* (also known as *GTL2*), *RTL1as* (*RTL1* antisense), and *MEG8*, and the differentially methylated region (DMR) of intergenic DMR and




Fig. 3. Fluorescence *in situ* hybridization using bacterial artificial chromosome clone probes RP11-35P13 (85,596,133–85,731,472) (green spectrum) at 14q31.3 and RP11-99C24 (89,927,464–90,094,722) (red spectrum) at 14q32.11. A direct duplication of 14q in the orientation of green–red–green–red is evident in the dup(14). The inset shows the amplified dup(14) and chromosome 14. dup(14) = the chromosome 14 with a duplication.

MEG3-DMR [19–22]. Our case did not involve duplication of the imprinting genes. However, our case did involve the duplication of other genes such as *FLRT2*, *GALC*, *GPR65*, *KCNK10*, *SPATA7*, *PTPN21*, *ZC3H14*, *EML5*, *TTC8*, *FOXN3*, *C14orf143*, *TDP1*, *KCNK13*, *PSMC1*, *C14orf102*, *CALM1*, *TTC7B*, *RPS6KA5*, *C14orf159*, *SNORA11B*, *GPR68*, and *CCDC88C*. The gene dosage increase effect of those genes is unknown at the present time. Although this reported neonate was normal in growth and psychomotor development with no apparent phenotypic abnormalities, long-term follow-ups are required for delineating the genotype–phenotype correlation of dup(14)(q31.3→q32.12).

Acknowledgments

This work was supported by research grants from the National Science Council (Grant Nos. NSC-99-2628-B-195-001-MY3 and NSC-101-2314-B-195-011-MY3) and from Mackay Memorial Hospital, Taipei, Taiwan (Grant No. MMH-E-101-04).

References

- [1] Orye E, Van Bever H, Desimpel H. Distal trisomy 14q due to tandem duplication (q24 leads to q32). *Ann Genet* 1983;26:238–9.
- [2] Turleau C, de Grouchy J, Chavin-Colin F, Denavit MF, Le Touzé P. Distal 14q trisomy. *Ann Genet* 1983;26:165–70 [Article in French].
- [3] Mikelsaar RV, Ilus TA, Lurie IW. Distal trisomy 14q. *J Med Genet* 1987;24:380–1.
- [4] Chen CP, Chern SR, Lin SP, Lin CC, Li YC, Wang TH, et al. A paternally derived inverted duplication of distal 14q with a terminal 14q deletion. *Am J Med Genet A* 2005;139A:146–50.
- [5] Thiel CT, Dörr HG, Trautmann U, Hoyer J, Hofmann K, Kraus C, et al. A *de novo* 7.6Mb tandem duplication of 14q32.2-qter associated with primordial short stature with neurosecretory growth hormone dysfunction, distinct facial anomalies and mild developmental delay. *Eur J Med Genet* 2008;51:362–7.
- [6] Trunca C, Opitz JM. Pericentric inversion of chromosome 14 and the risk of partial duplication of 14q (14q31 leads to 14qter). *Am J Med Genet* 1977;1:217–28.
- [7] Weinstein MR, Handmaker SD, Mohandas T. A unique pattern of dysmorphism and mental retardation associated with extra chromosome material (14q+): a probable duplication. *Am J Hum Genet* 1977;29:112A.
- [8] Sklower SL, Jenkins EC, Nolin SL, Duncan CJ, Warburton D, Yeboa KA, et al. Distal duplication 14q: report of three cases and further delineation of the syndrome. *Hum Genet* 1984;68:159–64.
- [9] Carr DM, Jones-Quartey K, Vartanian MV, Moore-Kaplan H. Duplication 14 (q31→qter). *J Med Genet* 1987;24:372–4.
- [10] Wakita Y, Narahara K, Kikkawa K, Namba H, Hiramoto K, Eguchi K, et al. Distal 14q trisomy syndrome in two siblings: further delineation of its phenotype. *Jinrui Idengaku Zasshi* 1988;33:469–75.
- [11] Masada CT, Olney AH, Fordyce R, Sanger WG. Partial deletion of 14q and partial duplication of 14q in sibs: testicular mosaicism for t(14q;14q) as a common mechanism. *Am J Med Genet* 1989;34:528–34.
- [12] Schinzel A. dup(14)(q31→qter). In: Schinzel A, editor. Catalogue of unbalanced chromosome aberrations in man. 2nd ed. Berlin: Walter de Gruyter, GmbH & Co.; 2001. p. 613–5.
- [13] Sonoda T, Kouno K, Sawada K, Sugimoto T. Distal 14q trisomy due to a maternal derivative chromosome 14. *Pediatr Int* 2001;43:313–5.
- [14] Chen CP, Kuo YK, Su YN, Chern SR, Tsai FJ, Wu PC, et al. Prenatal diagnosis and molecular cytogenetic characterization of a derivative chromosome der(18;18)(q10;q10)del(18)(q11.1q12.1) del(18)(q22.1q22.3) presenting as apparent isochromosome 18q in a fetus with holoprosencephaly. *Taiwan J Obstet Gynecol* 2011;50:182–7.
- [15] Chen CP, Ko TM, Su YN, Hsu CY, Chen YY, Su JW, et al. Prenatal diagnosis of partial trisomy 16p (16p12.2→pter) and partial monosomy 22q (22q13.31→qter) associated with increased nuchal translucency and abnormal maternal serum biochemistry in the first trimester. *Taiwan J Obstet Gynecol* 2012;51:129–33.
- [16] Chen CP, Su YN, Lin SP, Chern SR, Su JW, Chen YT, et al. Prenatal diagnosis and molecular cytogenetic characterization of a *de novo* interstitial duplication of 11q (11q22.3→q23.3) associated with abnormal maternal serum biochemistry. *Taiwan J Obstet Gynecol* 2013;52:120–4.
- [17] Kotzot D. Maternal uniparental disomy 14 dissection of the phenotype with respect to rare autosomal recessively inherited traits, trisomy mosaicism, and genomic imprinting. *Ann Genet* 2004;47:251–60.
- [18] Kotzot D. Prenatal testing for uniparental disomy: indications and clinical relevance. *Ultrasound Obstet Gynecol* 2008;31:100–5.
- [19] Hosoki K, Ogata T, Kagami M, Tanaka T, Saitoh S. Epimutation (hypomethylation) affecting the chromosome 14q32.2 imprinted region in a girl with upd(14)mat-like phenotype. *Eur J Hum Genet* 2008;16:1019–23.
- [20] Kagami M, Sekita Y, Nishimura G, Irie M, Kato F, Okada M, et al. Deletions and epimutations affecting the human 14q32.2 imprinted region in individuals with paternal and maternal upd(14)-like phenotypes. *Nat Genet* 2008;40:237–42.
- [21] Kagami M, O'Sullivan MJ, Green AJ, Watabe Y, Arisaka O, Masawa N, et al. The IG-DMR and the *MEG3*-DMR at human chromosome 14q32.2: hierarchical interaction and distinct functional properties as imprinting control centers. *PLoS Genet* 2010;6:e1000992.
- [22] Ogata T, Kagami M, Ferguson-Smith AC. Molecular mechanisms regulating phenotypic outcome in paternal and maternal uniparental disomy for chromosome 14. *Epigenetics* 2008;3:181–7.