




Contents lists available at ScienceDirect

Taiwanese Journal of Obstetrics & Gynecology

journal homepage: www.tjog-online.com

Research Letter

Molecular cytogenetic characterization and prenatal diagnosis of familial Xp22.33 microdeletion encompassing short stature homeobox gene in a male fetus with a favorable outcome



Chih-Ping Chen ^{a, b, c, d, e, f, *}, Tsang-Ming Ko ^g, Liang-Kai Wang ^a, Shuan-Pei Lin ^{b, h, i, j},
 Schu-Rern Chern ^b, Peih-Shan Wu ^k, Yen-Ni Chen ^a, Shin-Wen Chen ^a, Chien-Wen Yang ^b,
 Dai-Dyi Town ^a, Meng-Shan Lee ^a, Wayseen Wang ^{b, l}

^a Department of Obstetrics and Gynecology, MacKay Memorial Hospital, Taipei, Taiwan

^b Department of Medical Research, MacKay Memorial Hospital, Taipei, Taiwan

^c Department of Biotechnology, Asia University, Taichung, Taiwan

^d School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^e Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan

^f Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^g Genephile Bioscience Laboratory, Ko's Obstetrics and Gynecology, Taipei, Taiwan

^h Department of Pediatrics, MacKay Memorial Hospital, Taipei, Taiwan

ⁱ Department of Medicine, MacKay Medical College, New Taipei City, Taiwan

^j Department of Early Childhood Care, National Taipei University of Nursing and Health Sciences, Taipei, Taiwan

^k Gene Biodesign Co. Ltd, Taipei, Taiwan

^l Department of Bioengineering, Tatung University, Taipei, Taiwan

ARTICLE INFO

Article history:

Accepted 12 January 2017

Dear Editor,

A 34-year-old, gravida 1, para 0, woman underwent amniocentesis at 18 weeks of gestation because of advanced maternal age. She had a body height of 150 cm. Her husband was 38 years old and had a body height of 170 cm. The woman had a family history of short stature. Her elder sister had a body height of 148 cm, and her parents had a body height of 145 cm. However, her brother had a body height of 165 cm. Amniocentesis revealed a karyotype of 46,XY. Array comparative genomic hybridization (aCGH) analysis of amniotic fluid revealed a result of arr Xp22.33 (581,803–795,716) × 0–1 mat with a 213.9-kb Xp22.33 microdeletion encompassing only one Online Mendelian Inheritance in Man (OMIM) gene of short stature homeobox (*SHOX*). Multiplex ligation-dependent probe amplification-P018 (SALSA MLPA Probemix P018 *SHOX*; MRC-Holland, Amsterdam, The Netherlands)

analysis of the DNA extracted from the amniotic fluid and parental blood confirmed maternal transmission of the heterozygous *SHOX* deletion. The father did not have any *SHOX* deletion. The mother had a karyotype of 46,XX. An aCGH analysis of the DNA extracted from the maternal blood using CytoChip ISCA Array (Illumina, San Diego, CA, USA) revealed a result of arr Xp22.33 (553,160–1,232,886) × 1.3 with a 679.7-kb Xp22.33 microdeletion encompassing four genes including the OMIM gene of *SHOX* (Figure 1). The father had a karyotype of 46,XY. An aCGH analysis of the DNA extracted from the paternal blood revealed a result of arr (1–22) × 2, X × 1, Y × 1. The prenatal ultrasound findings were unremarkable. The parents elected to continue the pregnancy, and a normal male baby was delivered at 39 weeks of gestation with a body weight of 3126 g (>97th centile), body length of 49.5 cm (25–50th centile), head circumference of 32.5 cm (5–15th centile), and chest circumference of 31 cm. The cord blood had a karyotype of 46,XY. An aCGH analysis of the cord blood revealed a result of arr Xp22.33 (581,741–1,232,886) × 0.7 with a 651.1-kb Xp22.33 microdeletion encompassing four genes including the OMIM gene of *SHOX* (Figure 2). Metaphase fluorescence *in situ* hybridization analysis of 20 cultured lymphocytes obtained from the cord blood using the bacterial artificial chromosome probes RP11-808D8 (877,516–1,060,527) specific for Xpter (Xp22.33) crossing over with Ypter (Yp11.2) and RP11-943J20 (Xq11.1; 63,613,179–63,792,988) showed an Xp22.33 deletion, but no Yp11.2 deletion, in all 20 cells (Figure 3). Metaphase fluorescence *in situ* hybridization analysis of 20 cultured lymphocytes obtained from the maternal peripheral blood using

* Corresponding author. Department of Obstetrics and Gynecology, MacKay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, 10449, Taiwan.

E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).




Figure 1. Array comparative genomic hybridization analysis of the DNA extracted from maternal peripheral blood shows a 679.7-kb Xp22.33 microdeletion (553,160–1,232,886) encompassing *SHOX*. (A) Chromosome zoom-in view and (B) chromosome X. *SHOX* = short stature homeobox.

the bacterial artificial chromosome probes RP11-808D8 and RP11-943J20 showed a heterozygous Xp22.33 deletion in all 20 cells (Figure 4).

SHOX (OMIM 312865) (Xp22.33) encodes SHOX protein and is a pseudoautosomal homeobox-containing osteogenic gene. *SHOX* deficiency disorders include idiopathic familial short stature (OMIM 300582); Leri–Weill dyschondrosteosis (LWD) (OMIM 127300), which is inherited in a pseudoautosomal dominant pattern and has the classic clinical triad of short stature, mesomelia (the middle

portion of a limb is shortened in relation to the proximal portion), and Madelung deformity (abnormal alignment of the radius, ulna, and carpal bones at the wrist); and Langer mesomelic dysplasia (OMIM 249700), which is inherited in a pseudoautosomal recessive pattern resulted from *SHOX* nullizygosity due to biallelic *shox* deficiency and has severe short stature and severe skeletal dysplasia [1]. In case of LWD, short stature is noted in early childhood, mesomelia is evident in school-aged children, and Madelung deformity typically develops in mid-to-late childhood [1].




Figure 2. The aCGH analysis of the DNA extracted from cord blood shows a 651.1-kb Xp22.33 microdeletion (581,741–1,232,886) encompassing *SHOX*. (A) Chromosome zoom-in view and (B) chromosome X. aCGH = array comparative genomic hybridization; *SHOX* = short stature homeobox.

Differential diagnosis of isolated short stature and LWD caused by *SHOX* deficiency includes Turner syndrome in females, idiopathic short stature in both sexes, LWD caused by pathogenic variants other than *SHOX* deficiency, and trauma, infection, or tumors in the distal radial growth plate [1].

The peculiar aspect of the present case is the incidental detection of familial transmission of Xp22.33 microdeletion encompassing *SHOX* during prenatal genetic diagnosis because of

advanced maternal age. The present case provides evidence that prenatal diagnosis of Xp22.33 microdeletion encompassing *SHOX* can be associated with a favorable outcome. Recombinant human growth hormone therapy is effective in the treatment of patients with *SHOX* deficiency [1–18]. Prenatal diagnosis of familial *SHOX* deficiency provides the affected fetus an opportunity for early recognition of the disorder as well as early medical treatment in childhood.




Figure 3. Metaphase fluorescence *in situ* hybridization on the cord blood lymphocytes using the Xp22.33- and Yp11.2-specific probe of RP11-808D8 (Xp22.33 and Yp11.2; 877,516–1,060,527; fluorescein isothiocyanate, spectrum green) and the Xq11.1-specific probe of RP11-943J20 (Xq11.1; 63,613,179–63,792,988; Texas Red, spectrum red) shows one green signal in the Y chromosome and only one red signal in the del(X) chromosome with Xp deletion. del = deletion.




Figure 4. Metaphase FISH on the maternal peripheral blood lymphocytes using the Xp22.33- and Yp11.2-specific probe of RP11-808D8 (Xp22.33 and Yp11.2; 877,516–1,060,527; FITC, spectrum green) and the Xq11.1-specific probe of RP11-943J20 (Xq11.1; 63,613,179–63,792,988; Texas Red, spectrum red) shows one green signal and one red signal in the X chromosome and only one red signal in the del(X) chromosome with Xp deletion. del = deletion; FISH = fluorescence *in situ* hybridization; FITC = fluorescein isothiocyanate.

Conflicts of interest

The authors have no conflicts of interest relevant to this article.

Acknowledgments

This work was supported by research grants MOST-104-2314-B-195-009 and MOST-105-2314-B-195-012 from the Ministry of Science and Technology, and MMH-E-105-04 from MacKay Memorial Hospital, Taipei, Taiwan.

References

- [1] Binder G, Rappold GA. SHOX deficiency disorders. In: Pagon RA, Adam MP, Ardinger HH, Wallace SE, Amemiya A, Bean LJH, et al, editors. GeneReviews [Internet]. Seattle, WA: University of Washington; 1993–2016. Available at <https://www.ncbi.nlm.nih.gov/pubmed/20301394>. Accessed 2016 Dec 30.
- [2] Blum WF, Crowe BJ, Quigley CA, Jung H, Cao D, Ross JL, et al. Growth hormone is effective in treatment of short stature associated with short stature homeobox-containing gene deficiency: two-year results of a randomized, controlled, multicenter trial. *J Clin Endocrinol Metab* 2007;92:219–28.
- [3] Blum WF, Cao D, Hesse V, Fricke-Otto S, Ross JL, Jones C, et al. Height gains in response to growth hormone treatment to final height are similar in patients with SHOX deficiency and Turner syndrome. *Horm Res* 2009;71:167–72.
- [4] Blum WF, Ross JL, Zimmermann AG, Quigley CA, Child CJ, Kalifa G, et al. GH treatment to final height produces similar height gains in patients with SHOX deficiency and Turner syndrome: results of a multicenter trial. *J Clin Endocrinol Metab* 2013;98:E1383–92.
- [5] Shah BC, Moran ES, Zinn AR, Pappas JG. Effect of growth hormone therapy on severe short stature and skeletal deformities in a patient with combined Turner syndrome and Langer mesomelic dysplasia. *J Clin Endocrinol Metab* 2009;94:5028–33.
- [6] Cicognani A, Pirazzoli P, Nicoletti A, Baronio F, Conti V, Bonetti S. The SHOX gene: a new indication for GH treatment. *J Endocrinol Invest* 2010;33(Suppl. 6):15–8.
- [7] Iughetti L, Madeo S, Predieri B. Growth hormone therapy in patients with short stature homeobox-gene (SHOX) deficiency. *J Endocrinol Invest* 2010;33(Suppl. 6):34–8.
- [8] Iughetti L, Vannelli S, Street ME, Pirazzoli P, Bertelloni S, Radetti G, et al. Impaired GH secretion in patients with SHOX deficiency and efficacy of recombinant human GH therapy. *Horm Res Paediatr* 2012;78:279–87.
- [9] Lala R, Matarazzo P. SHOX mutation as a rare disease: molecular diagnosis and growth hormone treatment supported by the Italian public health system. *J Endocrinol Invest* 2010;33(Suppl. 6):39–43.
- [10] Richmond E, Rogol AD. Current indications for growth hormone therapy for children and adolescents. *Endocr Dev* 2010;18:92–108.
- [11] Scalco RC, Melo SSJ, Pugliese-Pires PN, Funari MFA, Nishi MY, Arnhold IJP, et al. Effectiveness of the combined recombinant human growth hormone and gonadotropin-releasing hormone analog therapy in pubertal patients with short stature due to SHOX deficiency. *J Clin Endocrinol Metab* 2010;95:328–32.
- [12] Takeda A, Cooper K, Bird A, Baxter L, Frampton GK, Gospodarevskaya E, et al. Recombinant human growth hormone for the treatment of growth disorders in children: a systematic review and economic evaluation. *Health Technol Assess* 2010;14. 1–209, iii–iv.
- [13] Binder G. Short stature due to SHOX deficiency: genotype, phenotype, and therapy. *Horm Res Paediatr* 2011;75:81–9.
- [14] Collett-Solberg PF. Update in growth hormone therapy of children. *J Clin Endocrinol Metab* 2011;96:573–9.
- [15] Kirk J. Indications for growth hormone therapy in children. *Arch Dis Child* 2012;97:63–8.
- [16] Loche S, Carta L, Ibba A, Guzzetti C. Growth hormone treatment in non-growth hormone-deficient children. *Ann Pediatr Endocrinol Metab* 2014;19:1–7.
- [17] Child CJ, Kalifa G, Jones C, Ross JL, Rappold GA, Quigley CA, et al. Radiological features in patients with short stature homeobox-containing (SHOX) gene deficiency and Turner syndrome before and after 2 Years of GH treatment. *Horm Res Paediatr* 2015;84:14–25.
- [18] Wit JM, Oostdijk W. Novel approaches to short stature therapy. *Best Pract Res Clin Endocrinol Metab* 2015;29:353–66.