



Contents lists available at ScienceDirect

## Taiwanese Journal of Obstetrics &amp; Gynecology

journal homepage: [www.tjog-online.com](http://www.tjog-online.com)

## Case Report

## Prenatal diagnosis and molecular cytogenetic characterization of mosaicism for a small supernumerary marker chromosome derived from chromosome 21q11.2-q21.1 and a literature review



Chih-Ping Chen <sup>a, b, c, d, e, f, \*</sup>, Ming Chen <sup>g, h, i</sup>, Chia-Hsun Wu <sup>a</sup>, Chen-Ju Lin <sup>a, j</sup>,  
 Schu-Rern Chern <sup>b</sup>, Peih-Shan Wu <sup>k</sup>, Yen-Ni Chen <sup>a</sup>, Shin-Wen Chen <sup>a</sup>,  
 Shun-Ping Chang <sup>g, h</sup>, Li-Feng Chen <sup>a</sup>, Wayseen Wang <sup>b, l</sup>

<sup>a</sup> Department of Obstetrics and Gynecology, MacKay Memorial Hospital, Taipei, Taiwan<sup>b</sup> Department of Medical Research, MacKay Memorial Hospital, Taipei, Taiwan<sup>c</sup> Department of Biotechnology, Asia University, Taichung, Taiwan<sup>d</sup> School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan<sup>e</sup> Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan<sup>f</sup> Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan<sup>g</sup> Department of Medical Research, Center for Medical Genetics, Changhua Christian Hospital, Changhua, Taiwan<sup>h</sup> Department of Genomic Medicine, Center for Medical Genetics, Changhua Christian Hospital, Changhua, Taiwan<sup>i</sup> Department of Obstetrics and Gynecology, Changhua Christian Hospital, Changhua, Taiwan<sup>j</sup> Department of Medicine, MacKay Medical College, Taipei, Taiwan<sup>k</sup> Gene Biodesign Co. Ltd, Taipei, Taiwan<sup>l</sup> Department of Bioengineering, Tatung University, Taipei, Taiwan

## ARTICLE INFO

## Article history:

Accepted 2 June 2017

## Keywords:

21q11.2-q21.1 duplication

Array comparative genomic hybridization

Chromosome 21

Small supernumerary marker chromosome

## ABSTRACT

**Objective:** We present prenatal diagnosis and molecular cytogenetic characterization of mosaicism for a small supernumerary marker chromosome (sSMC) derived from chromosome 21q11.2-q21.1, and we review the literature of an sSMC(21) with a duplication of 21q11.2-q21.1.

**Case report:** A 40-year-old woman underwent amniocentesis at 17 weeks of gestation because of advanced maternal age. Amniocentesis revealed a karyotype of 47,XX,+mar [18]/46,XX [4]. The parental karyotypes were normal. Prenatal ultrasound findings were unremarkable. aCGH analysis of cultured amniocytes revealed a 2.855-Mb duplication of 21q11.2-q21.1 encompassing the genes of *LIPI*, *ABCC13* and *NRIP1*. Metaphase fluorescence *in situ* hybridization analysis on cultured amniocytes revealed a result of 47,XX,+mar ish der(13/21) (D13/21Z1+) [10]. Spectral karyotyping analysis determined the origin of chromosome 21 in the sSMC. A female fetus was delivered with no phenotypic features of Down syndrome and no structural abnormalities. We discuss the genotype–phenotype correlation of *LIPI*, *ABCC13* and *NRIP1*, and review the literature of an sSMC(21) associated with dup(21)(q11.2q21.1).

**Conclusion:** aCGH is useful for identification of the nature and genetic component of a prenatally detected sSMC.

© 2017 Taiwan Association of Obstetrics & Gynecology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

A small supernumerary marker chromosome (sSMC) is an extra chromosome that has a size equal to or smaller than a chromosome 20 and cannot be identified by conventional cytogenetic technology

[1–3]. An sSMC derived from a non-acrocentric chromosome carries a higher risk for phenotypic abnormalities than an sSMC derived from an acrocentric chromosome (28% vs. 7%) [4]. We previously reported prenatal diagnosis and molecular cytogenetic characterization of an sSMC derived from chromosome 21 [5]. We previously also reported the application of array comparative genomic hybridization (aCGH) in identification of the nature and genetic component of an sSMC [6,7]. Here, we present an additional case of prenatal diagnosis of an sSMC(21) with euchromatic

\* Corresponding author. Department of Obstetrics and Gynecology, MacKay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei 10449, Taiwan.  
 E-mail address: [cpc\\_mmh@yahoo.com](mailto:cpc_mmh@yahoo.com) (C.-P. Chen).

material detected by aCGH. We also review the literature of an sSMC(21) with a duplication of chromosome 21q11.2-q21.1.

### Case report

A 40-year-old woman, gravid 2, para 1, underwent amniocentesis at 17 weeks of gestation because of advanced maternal age. Her husband was 44 years old. The woman and her husband were normal, and there was no family history of congenital malformations. Amniocentesis revealed a karyotype of 47,XX,+mar [18]/46,XX [4]. Among 22 colonies of cultured amniocytes, 18 colonies had a karyotype of 47,XX,+mar (Fig. 1), whereas the rest four colonies had a karyotype of 46,XX. The parental karyotypes were normal. Prenatal ultrasound findings were unremarkable. aCGH analysis of cultured amniocytes by SurePrint G3 Unrestricted CGH ISCA v2, 8 × 60 K Array (Agilent Technologies, Santa Clara, CA, USA) revealed a result of arr 21q11.2-q21.1 (15,485,008–18,340,054) × 3.0 with a 2.855-Mb duplication encompassing 10 Online Mendelian Inheritance of in Man (OMIM) genes of *LIPI*, *ABCC13*, *HSPA13*, *SMASN1*, *NRIP1*, *USP25*, *MIR99AHG*, *MIR99A*, *MIRLET7C* and *MIR125B2* (Fig. 2). Metaphase fluorescence *in situ* hybridization (FISH) analysis on cultured amniocytes revealed a result of 47,XX,+mar ish der(13/21)(D13/21Z1+) [10] (Fig. 3). Spectral karyotyping (SKY) analysis determined the origin of chromosome 21 in the sSMC (Fig. 4). The parents elected to terminate the pregnancy, and a female fetus was delivered with no phenotypic features of Down syndrome and no structural abnormalities. aCGH analysis of the DNA extracted from the skin revealed a result of arr 21q11.2-q21.1 (15,485,008–18,340,054) × 3.0. Cytogenetic analysis of the chorionic villi revealed a karyotype of 47,XX,+mar[27]/46,XX [13].

### Discussion

The present case had an sSMC(21) with a duplication of 21q11.2-q21.1 without involving Down syndrome critical region of 21q22.13-q22.3 and presented no Down syndrome phenotype.

Capkova et al. [8] reported a 4-year-old boy with hypotonia, brachycephaly, sandal gaps, joint hyperlaxity, developmental delay and speech delay with partial trisomy 21q11.2-q21.3, and a 6-year-old boy with macroglossia, hypotonia, joint hyperlaxity and normal mental development with partial tetrasomy 21q11.2 and partial trisomy 21q11.2-q21.3. To date, at least eight cases with an sSMC(21) and a partial duplication encompassing 21q11.2-q21.1 have been reported of which three cases are without clinical findings, and five cases are with clinical findings [2,9–14]. Liehr et al. [2] reported prenatal diagnosis of mosaicism for an sSMC(21) derived from 21p11.2-q21.1 and a phenotypically normal boy at one year of age. Liehr [14] reported a supernumerary r(21)(::p11.2 → q21.1::) in a normal female and her unborn child. Polityko et al. [10] reported a supernumerary r(21)(::p11.2 → q21.1::) in a normal male of whom the female partner had reported abortions. Liehr [14] reported 25% (5/20) mosaicism for an sSMC(21) derived from 21pter-q21.1 in the blood of a 62-year-old male with psychomotor retardation, severe developmental delay, no sexual development, short stature and microcephaly. Hamid et al. [13] and Liehr [14] reported 33.3% (5/15) mosaicism for an sSMC(21) derived from 21pter-q21.1 in the blood of a 4-year-old male with scoliosis, depressed nasal bridge, delayed fine motor skills and autism. Viersbach et al. [9] reported 70% mosaicism for a supernumerary r(21)(p11.2q21) in the blood of a 7-year-old female with facial dysmorphism, hypotonia and slightly developmental retardation. Baldwin et al. [11] reported 75% mosaicism for an sSMC(21) derive from 21pter → q21.1 in a 19-year-old individual with borderline normal intelligence, mild facial dysmorphism, bipolar disorder and diabetes mellitus. Liehr [14] reported 11.3% (6/53) mosaicism for an sSMC(21) derived from 21pter-q21.1 in the blood of an 8-year-old female with seizures, regression and mental retardation.

The present case had a duplication of 21q11.2. Niikawa et al. [15] and Shen et al. [16] mapped the putative transient abnormal myelopoiesis (TAM) gene at 21q11.2. TAM or transient myeloproliferative syndrome (OMIM 159595) is a leukemoid reaction that may occur in the newborn infants with Down syndrome [17]. The






Fig. 1. A karyotype of 47,XX,+mar. mar = marker chromosome.




**Fig. 2.** Array comparative genomic hybridization analysis on cultured amniocytes shows a 2.855-Mb duplication of 21q11.2-q21.1 encompassing the genes of *LIPI*, *ABCC13* and *NRIP1*. (A) and (B) chromosome 21 zoom-in views.

present case had a 2.855-Mb duplication of 21q11.2-q21.1 encompassing *LIPI*, *ABCC13* and *NRIP1*. Mutations of *LIPI* (OMIM 609252) are associated with autosomal dominant susceptibility to hypertriglyceridemia (OMIM 145750) [18]. Single nucleotide polymorphisms (SNPs) in *ABCC13* (OMIM 608835) have been associated with non-word repetition and reading measures [19]. *NRIP1* (OMIM 602490) is located at 21q11.2-q21.1 and encodes the transcriptional coregulator of receptor-interacting protein 140 (RIP140) or nuclear receptor-interacting protein 1 (NRIP1). NRIP1 or RIP140 is involved in the regulation of various oncogenic signaling pathways and participates in the development and progression of solid tumors and chronic lymphocytic leukemia [20]. Over-expression of NRIP1 is associated with psoriasis [21]. NRIP1 over-expression in Down syndrome down-regulates peroxisome proliferator-activated receptor  $\gamma$ , coactivator 1 $\alpha$  (PGC-1 $\alpha$  or *PPARGC1A*) and nuclear-encoded mitochondrial genes (NEMGs), and causes mitochondrial dysfunction in Down syndrome [22].

In summary, we present prenatal diagnosis and molecular cytogenetic characterization of mosaicism for an sSMC(21) derived from chromosome 21q11.2-q21.1. Our presentation demonstrates



**Fig. 3.** Metaphase fluorescence *in situ* hybridization analysis using the CEP13/21 FISH probe (D13/21Z1; spectrum green) and the CEP14/22 FISH probe (D14/22Z1; spectrum red) (Cytocell Inc, Adderbury, Oxfordshire, UK) shows four red signals and five green signals. The marker chromosome (arrow) presents a green signal indicating that the marker chromosome is derived from chromosome 13/21.



**Fig. 4.** The spectral karyotyping (SKY) analysis using 24-color SKY probes (Applied Spectral Imaging, Carlsbad, CA, USA) shows that the small supernumerary marker chromosome (arrow) is derived from chromosome 21.

the usefulness of aCGH in the identification of the nature and genetic component of a prenatally detected sSMC.

### Conflict of interest

The authors have no conflicts of interest relevant to this article.

### Acknowledgements

This work was supported by research grants MOST-104-2314-B-195-009 and MOST-105-2314-B-195-012 from the Ministry of Science and Technology and MMH-E-106-04 from MacKay Memorial Hospital, Taipei, Taiwan.

### References

- [1] Liehr T, Claussen U, Starke H. Small supernumerary marker chromosomes (sSMC) in humans. *Cytogenet Genome Res* 2004;107:55–67.
- [2] Liehr T, Mrasek K, Weise A, Dufke A, Rodríguez L, Martínez Guardia N, et al. Small supernumerary marker chromosomes – progress towards a genotype-phenotype correlation. *Cytogenet Genome Res* 2006;112:23–34.
- [3] Liehr T. Characterization of prenatally assessed *de novo* small supernumerary marker chromosomes by molecular cytogenetics. *Methods Mol Biol* 2008;444:27–38.
- [4] Crolla JA, Long F, Rivera H, Dennis NR. FISH and molecular study of autosomal supernumerary marker chromosomes excluding those derived from chromosomes 15 and 22: I. Results of 26 new cases. *Am J Med Genet* 1998;75:355–66.
- [5] Chen C-P, Lin C-C, Ko T-M, Tsai F-J, Chern S-R, Lee C-C, et al. Prenatal diagnosis and molecular cytogenetic characterization of a small supernumerary marker chromosome derived from chromosome 21. *Taiwan J Obstet Gynecol* 2010;49:377–80.
- [6] Chen C-P, Lin S-P, Lin Y-H, Chern S-R, Wu P-C, Chen Y-N, et al. Molecular cytogenetic characterization of mosaicism for a small supernumerary marker chromosome derived from chromosome 8 or r(8)(::p12→q13.1::) associated with phenotypic abnormalities. *Taiwan J Obstet Gynecol* 2016;55:852–5.
- [7] Chen C-P, Lin S-P, Chern S-R, Wu P-C, Chen Y-N, Chen S-W, et al. Molecular cytogenetic characterization of mosaicism for a small supernumerary marker chromosome derived from chromosome 8 or r(8)(::p11.22→q11.21::) in an 18-year-old female with short stature, obesity, attention deficit hyperactivity disorder and intellectual disability. *Taiwan J Obstet Gynecol* 2016;55:856–60.
- [8] Capkova P, Misovicova N, Vrbicka D. Partial trisomy and tetrasomy of chromosome 21 without down syndrome phenotype and short overview of genotype-phenotype correlation. A case report. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2014;158:321–5.
- [9] Viersbach R, Engels H, Gamberdinger U, Hansmann M. Delineation of supernumerary marker chromosomes in 38 patients. *Am J Med Genet* 1998;76:351–8.
- [10] Polityko AD, Rumyantseva NV, Starke H, Liehr T. Small supernumerary marker chromosomes characterized by multicolor FISH techniques: database of Belarusian national registry of chromosomal abnormalities. *Eur J Hum Genet* 2005;13(Suppl. 1):141–2 (Abstractno. P0289).
- [11] Baldwin EL, May LF, Justice AN, Martin CL, Ledbetter DH. Mechanisms and consequences of small supernumerary marker chromosomes: from Barbara McClintock to modern genetic-counseling issues. *Am J Hum Genet* 2008;82:398–410.
- [12] Manvelyan M, Riegel M, Santos M, Fuster C, Pellestor F, Mazauric M-L, et al. 32 new cases with small supernumerary marker chromosomes (sSMC) detected in connection with fertility problems – detailed molecular cytogenetic characterization and review of the literature. *Int J Mol Med* 2008;21:705–14.
- [13] Hamid AB, Kreskowski K, Weise A, Kosayakova N, Mrasek K, Voigt M, et al. How to narrow down chromosomal breakpoints in small and large derivative chromosomes – a new probe set. *J Appl Genet* 2012;53:259–69.
- [14] Liehr T. Small supernumerary marker chromosomes: chromosome 21. Available at: <http://ssmc-tl.com/chromosome-21.html>. [Accessed 15 March 2017].
- [15] Niikawa N, Deng H-X, Abe K, Harada N, Okada T, Tsuchiya H, et al. Possible mapping of the gene for transient myeloproliferative syndrome at 21q11.2. *Hum Genet* 1991;87:561–6.
- [16] Shen JJ, Williams BJ, Zipursky A, Doyle J, Sherman SL, Jacobs PA, et al. Cytogenetic and molecular studies of down syndrome individuals with leukemia. *Am J Hum Genet* 1995;56:915–25.
- [17] Seibel NL, Sommer A, Miser J. Transient neonatal leukemoid reactions in mosaic trisomy 21. *J Pediatr* 1984;104:251–4.
- [18] Wen XY, Hegele RA, Wang J, Wang DY, Cheung J, Wilson M, et al. Identification of a novel lipase gene mutated in *lpl* mice with hypertriglyceridemia and associated with dyslipidemia in humans. *Hum Mol Genet* 2003;12:1131–43.
- [19] Luciano M, Evans DM, Hansell NK, Medland SE, Montgomery GW, Martin NG, et al. A genome-wide association study for reading and language abilities in two population cohorts. *Genes Brain Behav* 2013;12:645–52.
- [20] Lapiere M, Castet-Nicolas A, Gitenay D, Jalaguier S, Teyssier C, Bret C, et al. Expression and role of RIP140/NRIP1 in chronic lymphocytic leukemia. *J Hematol Oncol* 2015;8:20.
- [21] Luan C, Chen X, Hu Y, Hao Z, Osland JM, Chen X, et al. Overexpression and potential roles of NRIP1 in psoriasis. *Oncotarget* 2016;7:74236–46.
- [22] Izzo A, Manco R, Bonfiglio F, Cali G, De Cristofaro T, Patergnani S, et al. NRIP1/RIP140 siRNA-mediated attenuation counteracts mitochondrial dysfunction in down syndrome. *Hum Mol Genet* 2014;23:4406–19.