



Research Letter

Fetal distress and urgent cesarean delivery due to new-onset peripartum Crohn's disease

Misgav Rottenstreich ^{a,*}, Amihai Rottenstreich ^b, Moshe Rottenstreich ^c^a Department of Obstetrics and Gynecology, Shaare Zedek Medical Center, Hebrew University Medical School of Jerusalem, Jerusalem, Israel^b Department of Obstetrics and Gynecology, Hadassah-Hebrew University Medical Center, Jerusalem, Israel^c Department of Gastroenterology, Shaare Zedek Medical Center, Hebrew University Medical School of Jerusalem, Jerusalem, Israel

ARTICLE INFO

Article history:

Accepted 1 August 2018

Dear Editor,

The incidence of inflammatory bowel diseases (IBDs) is increased among reproductive aged women [1]. The presence of active disease at early pregnancy is associated with an elevated risk for maternal and perinatal complications including low-birth weight infants, stillbirth and prematurity [3–5].

A 25 year-old Gravida-4 Para-3 presented at 36th weeks of gestation to the obstetric emergency room following the onset of uterine contractions lasting for few hours. She looked pale and weak with mild tachycardia (115 bpm), abdomen was soft with normal uterine tonus and vaginal examination revealed unfavorable cervix. Fetal heart rate monitoring demonstrated decreased variability with recurrent prolonged decelerations. She was immediately admitted to the operation room and an emergent cesarean section (CS) was performed. A 2674 g male was delivered with normal Apgar scores (9¹/9⁵). No apparent reason for fetal distress was observed intraoperatively, however, initial maternal laboratory tests on admission showed anemia of 7 g/dL with hematocrit of 22% WBC of $6 \times 10^3/\mu\text{L}$ blood glucose levels of 51 mg/dL and normal electrolytes. Following delivery, she was treated with two units of packed cells.

During hospitalization, she mentioned having watery-brown diarrhea up to 10 times/day lasting for 3 weeks prior to her admission. She denied abdominal pain, recent travelling, exposure to antibiotics or personal and family history of IBD. Repeat laboratory work-up revealed hypokalemia of 3.1 mEq/L, decreased serum albumin (1.9 g/dL) and markedly elevated C-reactive protein (39 mg/L, upper normal limit-0.5). WBC count increased up to




Fig. 1. CT scan showing large bowel distention (blue arrows) (A, B) and significant wall thickening (red arrows) (B, C).

* Corresponding author. Department of Obstetrics and Gynecology, Shaare Zedek Medical Center, 12 Bayit Street, 91031, Jerusalem, Israel. Fax: +972 2 666 6053.

E-mail address: misgavr@gmail.com (M. Rottenstreich).

$21.3 \times 10^3/\text{uL}$ at the fifth postoperative day. Stool cultures for bacteria and parasites, as well as *Clostridium Difficile* testing were negative. Thyroid function and serology for celiac disease and Rota virus were normal.

A computed tomography scan performed showed dilatation of the right and transverse colon and cecum up to 7.5 cm, with marked colitis from the splenic flexure through the rectum. The small intestine was otherwise normal (Fig. 1).

Sigmoidoscopy performed on the same day showed ulcerated, cobblestoned mucosa with scattered ulceration in the descending colon. Biopsy confirmed the diagnosis of CD. She was given intravenous corticosteroids along with total parental nutrition which led to resolution of her symptoms.

She was discharged 24 days postoperatively in good medical condition.

The occurrence of new-onset CD during pregnancy is rare [6–11]. Evaluation and diagnosis of suspected IBD during pregnancy is challenging since typical symptoms may be masked by normal physiologic changes coupled with difficulty in performing adequate radiologic and endoscopic evaluation during pregnancy due to safety considerations [12].

In our case, an emergent CS was performed due to fetal distress. This is in accordance with previous reports which demonstrated higher rates of CS, mostly unplanned, among IBD pregnant patients [5,13]. Fetal distress was reported as the leading indication [14]. As disease course may be fulminant, early diagnosis and prompt treatment is required due to high rate of maternal and perinatal complications.

Conflict of interest

All authors have no conflict of interest or benefit that has arisen from the direct applications this research to disclose.

Funding

None of the authors have any direct or indirect financial incentive associated with publishing the article.

Acknowledgements

Patient consent form has been completed and signed by the patient.

References

- [1] Cappell MS, Friedel D. Abdominal pain during pregnancy. *Gastroenterol Clin N Am* 2003;32:1–58.
- [2] Abhyankar A, Ham M, Moss AC. Meta-analysis: the impact of disease activity at conception on disease activity during pregnancy in patients with inflammatory bowel disease. *Aliment Pharmacol Ther* 2013;38:460–6. <https://doi.org/10.1111/apt.12417>.
- [3] Broms G, Granath F, Linder M, Stephansson O, Elmberg M, Kieler H. Birth outcomes in women with inflammatory bowel disease: effects of disease activity and drug exposure. *Inflamm Bowel Dis* 2014;20:1091–8. <https://doi.org/10.1097/MIB.0000000000000060>.
- [4] Cornish J, Tan E, Teare J, Teoh TG, Rai R, Clark SK, et al. A meta-analysis on the influence of inflammatory bowel disease on pregnancy. *Gut* 2007;56:830–7. <https://doi.org/10.1136/gut.2006.108324>.
- [5] Domeniconi D, Ferraris R, Manca A, Asnaghi G. Crohn disease of the colon with acute onset at the 17th week of pregnancy. Description of a clinical case. *Minerva Dietol Gastroenterol* 1989;35:135–7.
- [6] Seiner P, Olivero F, Maccario S, Libanori E, Meduri B, Ferrara L. Crohn's disease and pregnancy. Report of a clinical case with acute onset in the 33d week. *Minerva Ginecol* 1988;40:485–8.
- [7] Moeller DD. Crohn's disease beginning during pregnancy. *South Med J* 1988;81:1067.
- [8] Furderer T, Mantion G, Heyd B. Emergency surgery for intestinal obstruction revealing ileal Crohn's disease (CD) during pregnancy. *BMJ Case Rep* 2015;2015. <https://doi.org/10.1136/bcr-2013-009468>.
- [9] Zupi E, Valli E, Allegra E, Marconi D, Baschieri L. Crohn disease in pregnancy. Considerations on and discussion of a clinical case. *Minerva Ginecol* 1992;44:339–42.
- [10] Goettler CE, Stellato TA. Initial presentation of Crohn's disease in pregnancy: report of a case. *Dis Colon Rectum* 2003;46:406–10. <https://doi.org/10.1097/01.DCR.0000054697.05017.6C>.
- [11] Schulze H, Esters P, Dignass A. Review article: the management of Crohn's disease and ulcerative colitis during pregnancy and lactation. *Aliment Pharmacol Ther* 2014;40:991–1008. <https://doi.org/10.1111/apt.12949>.
- [12] Shand AW, Chen JS, Selby W, Solomon M, Roberts CL. Inflammatory bowel disease in pregnancy: a population-based study of prevalence and pregnancy outcomes. *BJOG An Int J Obstet Gynaecol* 2016;123:1862–70. <https://doi.org/10.1111/1471-0528.13946>.
- [13] Burke KE, Haviland MJ, Hacker MR, Shainker SA, Cheifetz AS. Indications for mode of delivery in pregnant women with inflammatory bowel disease. *Inflamm Bowel Dis* 2017;23:721–6. <https://doi.org/10.1097/MIB.0000000000001113>.